


REVIEW FOR FINAL

Remember that all exams are cumulative. I will likely recycle one or more questions from the previous exams. This review sheet is not meant to be exclusive. I will look at homework (including the mathematical part of computer homework), previous review sheets, previous exams, and Professor Stroyan's review material for ideas for the exam. You should too.

- (1) Find parametric equations for the line segment L from $A = (1, 4, 3)$ to $B = (2, 7, -4)$.
- (2) Find an equation for the plane P that is perpendicular to the line segment L in the previous exercise and which passes through the point $A = (1, 4, 3)$.
- (3) Find the distance from the point $C = (4, 42)$ to the plane in exercise 2. Here is how to do this: Consider the vector \vec{AC} from A to C . Project this vector on the line L , which is perpendicular to the plane. The length of the projection is the desired distance.
- (4) Let $f(x, y)$ be a function of two variables with continuous partial derivatives. What is the equation of the tangent plane to the graph of $z = f(x, y)$ at a point $(x_0, y_0, z_0 = f(x_0, y_0))$?
- (5) With the hypothesis of the previous exercise, what is the direction (given by a unit vector in the (x, y) -plane) of most rapid increase of the function f at the point (x_0, y_0) .
- (6) Continuing with the hypothesis of the previous exercise, what is a tangent vector to the graph of $z = f(x, y)$ at a point $(x_0, y_0, z_0 = f(x_0, y_0))$ which points in the direction of most rapid increase of the function f . Hint1: This was an e-exam. Hint2: the first two components of this 3D vector is the 2D vector of the previous exercise.
- (7) Redo the previous three exercises with the particular function $f(x, y) = x^2y + x$ and the point $(x_0, y_0) = (1, 2)$.
- (8) Find the directional derivative of $f(x, y) = x^2y + x$ at the point $(x_0, y_0) = (1, 2)$ in the direction of the vector $X = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$.
- (9) Find an equation for the tangent line to the curve $x^2 + y^4 = 5$ at the point $(1, \sqrt{2})$.

- (10) Find an equation for the tangent plane to the surface $x^2 + y^4 + z^2 = 6$ at the point $(1, \sqrt{2}, 1)$.
- (11) Accurately state a version of the implicit function theorem, as given in lecture. Hint: look at your lecture notes.
- (12) Let $f(x, y) = 3e^{-x^2-y^2} (x + 2y^2)$. Here are 3D and contour graphs of this function.



The partial derivatives of f are

$$\frac{\partial f}{\partial x}(x, y) = -3e^{-x^2-y^2} (-1 + 2x^2 + 4xy^2).$$

$$\frac{\partial f}{\partial y}(x, y) = -6e^{-x^2-y^2} y (-2 + x + 2y^2).$$

Mathematica gives the solutions to the equations

$$\frac{\partial f}{\partial x}(x, y) = 0, \quad \frac{\partial f}{\partial y}(x, y) = 0$$

as

$$\left\{ \left\{ x \rightarrow \frac{1}{4}, y \rightarrow \frac{-\sqrt{7}}{2} \right\}, \left\{ x \rightarrow \frac{1}{4}, y \rightarrow \frac{\sqrt{7}}{2} \right\}, \right. \\ \left. \left\{ y \rightarrow 0, x \rightarrow -\left(\frac{1}{\sqrt{2}}\right) \right\}, \left\{ y \rightarrow 0, x \rightarrow \frac{1}{\sqrt{2}} \right\} \right\}$$

Classify these critical points as maxima, minima, or neither.

- (13) Find the power series for $\frac{1}{1+x^2}$ by first writing down the series for $\frac{1}{1-x}$, using this to obtain the series for $\frac{1}{1+x}$ by making an appropriate substitution, and finally using this to obtain the desired series. Show that the radius of convergence of the series is 1 (by using the test involving ratios of successive coefficients). Find the series for $\arctan(x)$ by integrating term by term.

(14) Estimate the error if the series

$$\sum_{k=0}^{\infty} \frac{\sin(7kx)}{3^k}$$

is truncated after 100 terms. That is, estimate

$$\left| \sum_{k=101}^{\infty} \frac{\sin(7kx)}{3^k} \right|.$$

(15) Let C be smooth curve in the plane and $\mathbf{F}(x, y) = \begin{bmatrix} f(x, y) \\ g(x, y) \end{bmatrix}$ a vector field defined in a region containing the curve. What is the definition of $\int_C \mathbf{F} \cdot d\mathbf{x}$.

Answer: Let $\mathbf{X}(t) = \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$, $a \leq t \leq b$ be a smooth parametrization of C .

Then

$$\begin{aligned} \int_C \mathbf{F} \cdot d\mathbf{x} &= \int_a^b \mathbf{F}(\mathbf{X}(t)) \cdot \mathbf{X}'(t) dt \\ &= \int_a^b (f(x(t), y(t))x'(t) + g(x(t), y(t))y'(t)) dt. \end{aligned}$$

(16) Let C be smooth curve in the plane and $\omega = f(x, y)dx + g(x, y)dy$ a differential one-form defined in a region containing the curve. What is the definition of $\int_C \omega$? Answer: Let $\mathbf{X}(t) = \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$, $a \leq t \leq b$ be a smooth parametrization of C . Then

$$\int_C \omega = \int_a^b (f(x(t), y(t))x'(t) + g(x(t), y(t))y'(t)) dt.$$

(17) Calculate $\int_C \mathbf{F} \cdot d\mathbf{x}$, where C is the straight line path from $(-3, 2)$ to $(4, 5)$ and $\mathbf{F}(x, y) = \begin{bmatrix} x^2y \\ \sin(x) \end{bmatrix}$.

(18) Is the vector field $\mathbf{F}(x, y) = \begin{bmatrix} x^2y \\ \sin(x) \end{bmatrix}$ conservative?

(19) State the three forms of Green's theorem (neutral differential form form, curl form, divergence form).

(20) Let C be a simple closed curve in the plane enclosing a region D . What is the connection between $\int_C x dy$ and the area of D ?

4

(21) Find the power series expansions for $\frac{1}{1+x}$, $\frac{1}{1+x^2}$, $\frac{1}{\sqrt{1+x^2}}$, $\frac{1}{(1-x)^3}$, e^{-x^3} , $\sin(x^2)$.