

## 5.5 Series Solutions Near a Regular Singular Point, Part I

Theorem 5.3.1: If  $p(x)$  and  $q(x)$  are analytic at  $x_0$  (i.e.,  $x_0$  is an ordinary point of the ODE  $y'' + p(x)y' + q(x)y = 0$ ), then the general solution to this ODE is

$$y = \sum_{n=0}^{\infty} a_n (x - x_0)^n = a_0 \phi_0(x) + a_1 \phi_1(x)$$

where  $\phi_i$  are power series solutions that are analytic at  $x_0$ . The solutions  $\phi_0, \phi_1$  form a fundamental set of solutions. The radius of convergence for each of these series solutions is at least as large as the minimum radii of convergence of the series for  $\frac{Q}{P}$  and  $\frac{R}{P}$ .

If you prefer a power series expansion about 0, use  $u$ -substitution: let  $u = x - x_0$ . Then  $p(u + x_0)$  and  $q(u + x_0)$  are analytic at 0

(Semi-failed) attempt to transform 5.5 problem into 5.4 problem:

**5.5:**  $y'' + p(x)y' + q(x)y = 0$

$$x^2 y'' + x^2 p(x) y' + x^2 q(x) y = 0$$

$x^2 y'' + x[xp(x)]y' + [x^2 q(x)]y = 0$  where  $xp(x)$  and  $x^2 q(x)$  are functions of  $x$ .

**5.4:**  $x^2 y'' + \alpha x y' + \beta y = 0$  where  $\alpha, \beta$  are constants.

Combine 5.3/5.4 methods.

Defn:  $x_0$  is a *regular singular value* if  $x_0$  is a singular value and  $xp(x)$  and  $x^2 q(x)$  are analytic at  $x_0$ . A singular value which is not regular is called *irregular*.

Examples:

$$y'' + \frac{y'}{x} + \frac{y}{x^2} = 0, \text{ regular singular value: } x = 0.$$

$$y'' + \frac{y'}{x^2} + \frac{y}{x} = 0, \text{ irregular singular value: } x = 0.$$

$$y'' + y' + \frac{y}{x^3} = 0, \text{ irregular singular value: } x = 0.$$

If  $p(x)$  and  $q(x)$  are rational functions, then  $xp(x)$  and  $x^2q(x)$  are analytic iff  $\lim_{x \rightarrow 0} xp(x)$  and iff  $\lim_{x \rightarrow 0} x^2q(x)$  are finite. (i.e., after reducing fractions,  $x$  is not in the denominator.

Ex:  $p(x) = \frac{1}{x}$  implies  $xp(x) = \frac{x}{x} = 1$

Ex:  $p(x) = \frac{1}{x^2}$  implies  $xp(x) = \frac{x}{x^2} = \frac{1}{x}$

---

If  $x_0 = 0$  is a regular singular value of the linear homogeneous DE,  $x^2y'' + x[xp(x)]y' + x^2q(x)y = 0$  (\*), then

$xp(x) = \sum_{n=0}^{\infty} p_n x^n$  and  $x^2q(x) = \sum_{n=0}^{\infty} q_n x^n$  for constants  $p_n, q_n$ .

If  $y = x^r \sum_{n=0}^{\infty} a_n x^n = \sum_{n=0}^{\infty} a_n x^{n+r}$  is a solution to (\*) where  $r \neq 0$ .

$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$  and  $y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$

$$x^2 \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} + x[xp(x)] \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} + [x^2q(x)] \sum_{n=0}^{\infty} a_n x^{n+r}$$

$$\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r} + [xp(x)] \sum_{n=0}^{\infty} (n+r) a_n x^{n+r} + [x^2q(x)] \sum_{n=0}^{\infty} a_n x^{n+r}$$

$$\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r} + (\sum_{n=0}^{\infty} p_n x^n) (\sum_{n=0}^{\infty} (n+r) a_n x^{n+r}) + (\sum_{n=0}^{\infty} q_n x^n) (\sum_{n=0}^{\infty} a_n x^{n+r})$$

Thus the coefficient of  $x^r$  is  $r(r-1)a_0 + p_0 r a_0 + q_0 a_0 = 0$

We can take  $a_0 \neq 0$ . Thus  $r(r-1) + p_0 r + q_0 = 0$

Thus we can solve for  $r$  using the quadratic formula.

Case 1:  $r_1 > r_2$  both real and  $r_1 - r_2$  is not an integer.

Case 2:  $r_1 > r_2$  both real and  $r_1 - r_2 = p$ ,  $p$  an integer.

Case 3: one repeated root.

Case 4: two complex roots.