


[3] 1.) Write the following in the form $u(t) = R\cos(bt - \delta)$

$$u(t) = 2\sqrt{3}\cos(2t) - 2\sin(2t) = \underline{\hspace{10cm}}$$



[3] 2.) If $y = \phi_1(t)$ and $y = \phi_2(t)$ are homogeneous solutions to $ay'' + by' + cy = g(t)$, then a non-homogeneous solution to this second order linear differential equation is

$$y(t) = \underline{\hspace{10cm}}$$

where $u_1(t) = \underline{\hspace{10cm}}$ and $u_2(t) = \underline{\hspace{10cm}}$

[4] 3.) A spring mass system has a spring constant of 3N/m. A mass of 4kg is attached to the spring. The system is driven by an external force of $\sin(3t)$ N. The spring is stretched 5 m and then set in motion with a downward velocity of 2m/s. Assume that there is no damping. State the initial value problem that describes the motion of this mass. **Do NOT solve.**

IVP: $\underline{\hspace{10cm}}$

[10] 4.) Solve $y'' - 4y = -12 + 8e^{2t}$, $y(0) = 3$, $y'(0) = 6$.


Answer: _____

[10] 4.) Solve $y'' + 4y = 8 + \sin(2t)$, $y(0) = 4$, $y'(0) = 1$.

Answer: _____

[3] 1.) Write the following in the form $u(t) = R\cos(bt - \delta)$

$u(t) = -2\cos(2t) + 2\sqrt{3}\sin(2t) =$ _____



[3] 2.) If $y = \phi_1(t)$ and $y = \phi_2(t)$ are homogeneous solutions to $ay'' + by' + cy = g(t)$, then a non-homogeneous solution to this second order linear differential equation is

$y(t) =$ _____

where $u_1(t) =$ _____ and $u_2(t) =$ _____

[4] 3.) A spring is stretched 1m by a force of 2N. A mass of 9kg is attached to the spring and also attached to a viscous damper that exerts a force of 3N when the velocity of the mass is 1m/sec. The spring is set in motion from its equilibrium position with an upward velocity of 2m/s. State the initial value problem that describes the motion of this mass. **Do NOT solve.**

IVP: _____

[10] 4.) Solve $y'' - y = t + 24e^{3t}$, $y(0) = 5$, $y'(0) = 2$.

Answer: _____


[10] 4.) Solve $y'' + 2y' + 5y = 50 + 34\sin(2t)$, $y(0) = 4$, $y'(0) = 1$.

Answer: _____

[3] 1.) Write the following in the form $u(t) = R\cos(bt - \delta)$

$$u(t) = -2\cos(2t) + 2\sqrt{3}\sin(2t) = \underline{4\cos(2t - \frac{2\pi}{3})}$$


$$\frac{\pi}{2} + \frac{\pi}{6} = \frac{4\pi}{6} = \frac{2\pi}{3}$$



[3] 1.) Write the following in the form $u(t) = R\cos(bt - \delta)$


$$u(t) = -2\cos(2t) - 2\sqrt{3}\sin(2t) = \underline{4\cos(2t + \frac{2\pi}{3})}$$

$$-\frac{\pi}{2} - \frac{\pi}{6} = -\frac{4\pi}{6} = -\frac{2\pi}{3}$$




[3] 1.) Write the following in the form $u(t) = R\cos(bt - \delta)$

$$u(t) = 2\cos(2t) - 2\sqrt{3}\sin(2t) = \underline{4\cos(2t + \frac{\pi}{3})}$$



[3] 1.) Write the following in the form $u(t) = R\cos(bt - \delta)$

$$u(t) = 2\sqrt{3}\cos(2t) - 2\sin(2t) = \underline{4\cos(2t + \frac{\pi}{6})}$$



[3] 2.) If $y = \phi_1(t)$ and $y = \phi_2(t)$ are homogeneous solutions to $ay'' + by' + cy = g(t)$, then a non-homogeneous solution to this second order linear differential equation is

$$y(t) = u_1(t)\phi_1 + u_2(t)\phi_2$$

$$\text{where } u_1(t) = \int \frac{\begin{vmatrix} 0 & \phi_2 \\ 1 & \phi_2' \end{vmatrix}}{\begin{vmatrix} \phi_1 & \phi_2 \\ \phi_1' & \phi_2' \end{vmatrix}} \frac{g(t)}{a} dt \quad \text{and} \quad u_2(t) = \int \frac{\begin{vmatrix} \phi_1 & 0 \\ \phi_1' & 1 \end{vmatrix}}{\begin{vmatrix} \phi_1 & \phi_2 \\ \phi_1' & \phi_2' \end{vmatrix}} \frac{g(t)}{a} dt$$

[4] 3.) A 10 kg mass stretches a spring 5 m. The mass is acted on by an external force of $6\cos(3t)$ N. The spring is compressed 4 m and then set in motion with an upward velocity of 2m/s. Assume that there is no damping. State the initial value problem that describes the motion of this mass. **Do NOT solve.**

$$\text{IVP: } \underline{10u'' + 19.6u = 6\cos(3t), \quad u(0) = -4, u'(0) = -2}$$

$$mg = kL, \quad k = \frac{10(9.8)}{5} = 2(9.8) = 19.6$$

[4] 3.) A 20 kg mass stretches a spring 2 m. The mass is acted on by an external force of $9\sin(5t)$ N and moves in a medium that imparts of viscous force of 8N when the speed of the mass is 4m/sec. The spring is compressed 4 m and then released. State the initial value problem that describes the motion of this mass. **Do NOT solve.**

$$\text{IVP: } \underline{20u'' + 2u' + 98u = 9\sin(5t), \quad u(0) = -4, u'(0) = 0}$$

$$mg = kL, \quad k = \frac{20(9.8)}{2} = 98$$

$$8 = \gamma(4). \quad \text{Thus } \gamma = 2$$

[4] 3.) A spring mass system has a spring constant of 3N/m. A mass of 4kg is attached to the spring. The system is driven by an external force of $\sin(3t)$ N. The spring is stretched 5 m and then set in motion with a downward velocity of 2m/s. Assume that there is no damping. State the initial value problem that describes the motion of this mass. **Do NOT solve.**

$$\text{IVP: } \underline{4u'' + 3u = \sin(3t), \quad u(0) = 5, u'(0) = 2}$$

[4] 3.) A spring is stretched 1m by a force of 2N. A mass of 9kg is attached to the spring and also attached to a viscous damper that exerts a force of 3N when the velocity of the mass is 1m/sec. The spring is set in motion from its equilibrium position with an upward velocity of 2m/s. State the initial value problem that describes the motion of this mass. **Do NOT solve.**

$$\text{IVP: } \underline{9u'' + 3u' + 2u = 0, \quad u(0) = 0, u'(0) = -2}$$

$$2 = 1k. \quad \text{Thus } k = 2$$

$$3 = \gamma(1).$$

[10] 4.) Solve $y'' - 4y = -12 + 8e^{2t}$, $y(0) = 3$, $y'(0) = 6$.

Step 1: Solve homogeneous $y'' - 4y = 0$

Guess $y = e^{rt}$: $r^2 - 4 = 0$, Thus $r = \pm 2$

Homogeneous solution: $y = c_1e^{2t} + c_2e^{-2t}$

Step 2a: Find a non-homogeneous solution to $y'' - 4y = -12$

Guess $y = A$. Then $y' = 0$ and $y'' = 0$.

Plug in: $y'' - 4y = -12$: $0 - 4A = -12$.

Thus $A = 3$

Step 2b: Find a non-homogeneous solution to $y'' - 4y = 8e^{2t}$

Guess $y = Ate^{2t}$. Then $y' = Ae^{2t} + 2Ate^{2t}$ and $y'' = 2Ae^{2t} + 2Ae^{2t} + 4Ate^{2t} = 4Ae^{2t} + 4Ate^{2t}$.

Plug in: $y'' - 4y = 8e^{2t}$: $4Ae^{2t} + 4Ate^{2t} - 4Ate^{2t} = 8e^{2t}$

$4Ae^{2t} = 8e^{2t}$. Thus $A = 2$.

General solution: $y = c_1e^{2t} + c_2e^{-2t} + 3 + 2te^{2t}$

$y' = 2c_1e^{2t} - 2c_2e^{-2t} + 2(e^{2t} + 2te^{2t})$

Initial values: $y(0) = 3$, $y'(0) = 6$

$y(0) = 3$: $3 = c_1 + c_2 + 3$. Thus $c_2 = -c_1$

$y'(0) = 6$: $6 = 2c_1 - 2c_2 + 2$. Thus $4 = 2c_1 + 2c_1$. Thus $4 = 4c_1$.

Hence $c_1 = 1$ and $c_2 = -c_1 = -1$

IVP solution: $y = e^{2t} - e^{-2t} + 3 + 2te^{2t}$

Answer: $y = e^{2t} - e^{-2t} + 3 + 2te^{2t}$

[10] 4.) Solve $y'' - y = t + 24e^{3t}$, $y(0) = 5$, $y'(0) = 2$.

Step 1: Solve homogeneous $y'' - y = 0$

Guess $y = e^{rt}$: $r^2 - 1 = 0$, Thus $r = \pm 1$

Homogeneous solution: $y = c_1e^t + c_2e^{-t}$

Step 2a: Find a non-homogeneous solution to $y'' - y = t$

Guess $y = At + B$. Then $y' = A$ and $y'' = 0$.

Plug in: $-(At + B) = t$: $-A = 1$ and $B = 0$.

So non-homogeneous solution to $y'' - y = t$ is $y = -t$

Sidenote: No y' term means we did not need constant term B , but including it is fine.

Step 2b: Find a non-homogeneous solution to $y'' - y = 24e^{3t}$

Guess $y = Ae^{3t}$. Then $y' = 3Ae^{3t}$ and $y'' = 9Ae^{3t}$.

Plug in: $y'' - y = 24e^{3t}$: $9Ae^{3t} - Ae^{3t} = 24e^{3t}$ Thus $8Ae^{3t} = 24e^{3t}$ and $A = 3$.

General solution: $y = c_1e^t + c_2e^{-t} - t + 3e^{3t}$

$y' = c_1e^t - c_2e^{-t} - 1 + 9e^{3t}$

Initial values: $y(0) = 5$, $y'(0) = 2$

$y(0) = 5$: $5 = c_1 + c_2 - 0 + 3$. Thus $c_1 + c_2 = 2$

$y'(0) = 2$: $2 = c_1 - c_2 - 1 + 9$. Thus $c_1 - c_2 = -6$.

Thus $2c_1 = -4$ and thus $c_1 = -2$. Also $2c_2 = 8$ and $c_2 = 4$

IVP solution: $y = -2e^t + 4e^{-t} - t + 3e^{3t}$

Answer: $y = -2e^t + 4e^{-t} - t + 3e^{3t}$

[10] 4.) Solve $y'' + 2y' + 5y = 50 + 34\sin(2t)$, $y(0) = 4$, $y'(0) = 1$.

Step 1: Solve homogeneous DE: $y'' + 2y' + 5y = 0$

Guess $y = e^{rt}$: $r^2 + 2r + 5 = 0$, Thus $r = \frac{-2 \pm \sqrt{2^2 - 4(5)}}{2} = \frac{-2 \pm \sqrt{-16}}{2} = \frac{-2 \pm 4i}{2} = -1 \pm 2i$

Homogeneous solution: $y = c_1 e^{-t} \cos(2t) + c_2 e^{-t} \sin(2t)$

Step 2a: Find a non-homogeneous solution to $y'' + 2y' + 5y = 50$

Guess $y = A$. Then $y' = 0$ and $y'' = 0$.

Plug in: $y'' + 2y' + 5y = 50$: $0 + 5A = 50$. Thus $A = 10$

Step 2b: Find a non-homogeneous solution to $y'' + 2y' + 5y = 34\sin(2t)$

Guess $y = A\cos(2t) + B\sin(2t)$.

Then $y' = -2A\sin(2t) + 2B\cos(2t)$ and $y'' = -4A\cos(2t) - 4B\sin(2t)$.

Plug in: $y'' + 2y' + 5y = 34\sin(2t)$:

$$-4A\cos(2t) - 4B\sin(2t) + 2(-2A\sin(2t) + 2B\cos(2t)) + 5(A\cos(2t) + B\sin(2t)) = 34\sin(2t)$$

$$-4A\cos(2t) + 4B\cos(2t) + 5A\cos(2t) = 0. \text{ Thus } -4A + 4B + 5A = 0. \text{ Thus } A = -4B$$

$$-4B\sin(2t) - 4A\sin(2t) + 5B\sin(2t) = 34\sin(2t).$$

$$\text{Thus } -4B - 4A + 5B = 1. \text{ Thus } B - 4A = B - 4(-4B) = 17B = 34. \text{ Thus } B = 2 \text{ and } A = -8$$

General solution: $y = c_1 e^{-t} \cos(2t) + c_2 e^{-t} \sin(2t) + 10 - 8\cos(2t) + 2\sin(2t)$

$$y' = c_1[-e^{-t} \cos(2t) - 2e^{-t} \sin(2t)] + c_2[-e^{-t} \sin(2t) + 2e^{-t} \cos(2t)] + 16\sin(2t) + 4\cos(2t)$$

Initial values: $y(0) = 4$, $y'(0) = 1$.

$$y(0) = 4: \quad 4 = c_1 + 10 - 8. \text{ Thus } c_1 = 2$$

$$y'(0) = 1: \quad 1 = c_1[-1] + c_2[2] + 4. \text{ Thus } -2 = -2 + 2c_2. \text{ Thus } c_2 = 0$$

IVP solution: $y = 2e^{-t} \cos(2t) + 10 - 8\cos(2t) + 2\sin(2t)$

$$\text{Answer: } \underline{y = 2e^{-t} \cos(2t) + 10 - 8\cos(2t) + 2\sin(2t)}$$

[10] 4.) Solve $y'' + 4y = 8 + \sin(2t)$, $y(0) = 4$, $y'(0) = 1$.

Step 1: Solve homogeneous $y'' + 4y = 0$

Guess $y = e^{rt}$: $r^2 + 4 = 0$, Thus $r = \pm 2i$

Homogeneous solution: $y = c_1 \cos(2t) + c_2 \sin(2t)$

Step 2a: Find a non-homogeneous solution to $y'' + 4y = 8$

Guess $y = A$. Then $y' = 0$ and $y'' = 0$.

Plug in: $y'' + 4y = 8$: $0 + 4A = 8$. Thus $A = 2$

Step 2b: Find a non-homogeneous solution to $y'' + 4y = \sin(2t)$

Guess $y = t(A \cos(2t) + B \sin(2t))$.

Then $y' = t[-2A \sin(2t) + 2B \cos(2t)] + A \cos(2t) + B \sin(2t)$ and

$y'' = t[-4A \cos(2t) - 4B \sin(2t)] + -2A \sin(2t) + 2B \cos(2t) + -2A \sin(2t) + 2B \cos(2t)$.

Thus $y'' = (4B - 4At) \cos(2t) + (-4Bt - 4A) \sin(2t)$

Plug in: $y'' + 4y = \sin(2t)$:

$(4B - 4At) \cos(2t) + (-4Bt - 4A) \sin(2t) + 4[t(A \cos(2t) + B \sin(2t))] = \sin(2t)$

$(4B - 4At + 4tA) \cos(2t) + (-4Bt - 4A + 4tB) \sin(2t) = \sin(2t)$

$(4B) \cos(2t) + (-4A) \sin(2t) = \sin(2t)$

Thus $4B = 0$ and $B = 0$. Also $-4A = 1$ Thus $A = -\frac{1}{4}$

General solution: $y = c_1 \cos(2t) + c_2 \sin(2t) + 2 - \frac{1}{4}t \cos(2t)$

$y' = -2c_1 \sin(2t) + 2c_2 \cos(2t) - \frac{1}{4} \cos(2t) + \frac{1}{2}t \sin(2t)$

Initial values: $y(0) = 4$, $y'(0) = 1$

$y(0) = 4$: $4 = c_1 + 2$. Thus $c_1 = 2$

$y'(0) = 1$: $1 = 2c_2 - \frac{1}{4}$. Thus $c_2 = \frac{1}{2} + \frac{1}{8} = \frac{5}{8}$.

Answer: $y = 2 \cos(2t) + \frac{5}{8} \sin(2t) + 2 - \frac{1}{4}t \cos(2t)$