


A very elementary introduction to proofs

Part 2

Example: Prove a function is not 1:1



By Dr. Isabel Darcy,
Dept of Mathematics and AMCS,
University of Iowa

Some notation:

\forall = for all

\sim = negation

\exists = there exists

! = unique

$\exists !$ = there exists a unique

TFAE = The following are equivalent

$[p \Rightarrow q]$ is equivalent to $[\forall p, q \text{ holds}]$.

That is, for everything satisfying the hypothesis p , the conclusion q must hold.

$f : A \rightarrow B$ is 1:1 iff

$f(x_1) = f(x_2)$ implies $x_1 = x_2$.

$f : A \rightarrow B$ is 1:1 iff

$\forall x_1$ and $\forall x_2$ such that $f(x_1) = f(x_2)$,
we have $x_1 = x_2$.

How do we prove a function is **NOT** 1:1

A statement is **false** if the hypothesis holds, but the conclusion need not hold.

TFAE (The following are equivalent):

Hypothesis does not implies conclusion

p does not imply q.

$$p \not\Rightarrow q.$$

It is not true that $\forall p, q$ holds.

$\exists p$ such that q does not hold.

That is there exists a specific case where the hypothesis holds, but the conclusion does not hold.

$\sim [p \Rightarrow q]$ is equivalent to $\sim [\forall p, q \text{ holds}]$.

Thus if $p \Rightarrow q$ is false,

then it is not true that $[\forall p, q \text{ holds}]$.

That is, $\exists p$ such that q does not hold.

To prove that a statement is false:

Find an example where the hypothesis holds, but the conclusion does not hold.

$f : A \rightarrow B$ is 1:1 iff

$\forall x_1$ and $\forall x_2$ such that $f(x_1) = f(x_2)$,


we have $x_1 = x_2$.

Ex: To prove a function is not 1:1, find specific
 x_1, x_2 such that $f(x_1) = f(x_2)$, but $x_1 \neq x_2$.

Ex: To prove a function is not 1:1, find specific x_1, x_2 such that $f(x_1) = f(x_2)$, but $x_1 \neq x_2$.

Ex: $f: \mathbb{R} \rightarrow \mathbb{R}, f(x) = x^2$ is not 1:1

Proof:



$$f(1) = 1^2 = 1 = (-1)^2 = f(-1), \text{ but } 1 \neq -1$$

$$\rightarrow (-x)^2 = x^2$$

Does there exist x
st $-x \neq x$

Contrapositive of $[p \implies q]$ is $[\sim q \implies \sim p]$.

Example: The contrapositive of

$f(x_1) = f(x_2)$ implies $x_1 = x_2$.

is

$x_1 \neq x_2$ implies $f(x_1) \neq f(x_2)$.

Example: The contrapositive of

$\ln(x_1) = \ln(x_2)$ implies $x_1 = x_2$.

is

$x_1 \neq x_2$ implies $\ln(x_1) \neq \ln(x_2)$.

The contrapositive of a theorem is true:

If $p \Rightarrow q$ is true, then
its contrapositive $\sim q \Rightarrow \sim p$ is also true.

If the ^{$\sim q$} conclusion q does not hold,
then the hypothesis p cannot hold.
 _{$\sim p$}

If $x_1 \neq x_2$ then $\ln(x_1) \neq \ln(x_2)$,
since $f(x) = \ln(x)$ is 1:1.

Sidenote: The converse of $[p \Rightarrow q]$ is $[q \Rightarrow p]$.

The converse of a theorem need not be true.

That is $p \Rightarrow q$ does not imply $q \Rightarrow p$.

TFAE:

- ▶ $f : A \rightarrow B$ is 1:1.
- ▶ $f(x_1) = f(x_2)$ implies $x_1 = x_2$.
- ▶ $x_1 \neq x_2$ implies $f(x_1) \neq f(x_2)$.
- ▶ $\forall x_1 \neq x_2, f(x_1) \neq f(x_2)$.

$f : A \rightarrow B$ is NOT 1:1 iff $\exists \equiv$ Find $x_1 \neq x_2$

$\exists x_1 \neq x_2$ such that $f(x_1) = f(x_2)$.