

EX: 231

### 6.3 Derangements

Suppose each person in a group of  $n$  friends brings a gift to a party. In how many ways can the  $n$  gifts be distributed so that each person receives one gift and no person receives their own gift.

*Secret Santa*

Let the set of friends =  $\{p_1, \dots, p_n\}$  where  $p_j$  = person  $j$ .

Let the set of gifts =  $\{g_1, \dots, g_n\}$  where  $g_j$  = the gift brought by person  $j$ .

Suppose  $f : \{p_1, \dots, p_n\} \rightarrow \{g_1, \dots, g_n\}$ ,  $f(p_k) = g_j$  iff person  $p_k$  receives gift  $g_j$ , the gift brought by person  $j$ .


If each person receives one gift, then  $f$  is a bijection.

If no person receives their own gift. Then  $f(p_j) \neq g_j$ .

In simpler notation,  $f : \{1, \dots, n\} \rightarrow \{1, \dots, n\}$  such that  $f(j) \neq j$

Recall:

a permutation on  $\{1, \dots, n\}$  is a bijection  $f : \{1, \dots, n\} \rightarrow \{1, \dots, n\}$



Ex: The permutation 1 2 3 4 5 corresponds to the identity function.

Ex: The permutation 1 3 2 corresponds to the function  $f(1) = 1, f(2) = 3, f(3) = 2$


Defn: A derangement of  $\{1, \dots, n\}$  is a permutation  $i_1 i_2 \dots i_n$  such that  $i_j \neq j$ . I.e,  $j$  is not in the  $j$ th place.

In function notation:

$$f(j) = i_j, \text{ then if } i_1 i_2 \dots i_n \text{ is a derangement, } f(j) \neq j.$$

In yet other wording, recall a permutation corresponds to the placement of  $n$  non-attacking rooks on an  $n \times n$  chessboard.


Ex: The permutation 1 3 2 corresponds to the following rook placement:



3  
2  
1  
1 3 2  
nota derangement

A derangement corresponds to non-attacking rook placement with forbidden positions along the diagonal  $(j, j)$ , for  $j = 1, \dots, n$ .

Ex: If rooks are placed on the following  $3 \times 3$  chessboard in non-attacking position, then the rook placement corresponds to a derangement if no rook is placed in a spot marked with an X.




Thus the derangements of  $\{1, 2, 3\}$  are 2 3 1 and 3 1 2.

Let  $D_n$  = the number of derangements of  $\{1, \dots, n\}$ .

Thus  $D_3 = 2$ .

$$D_2 = 1$$



**Thm 6.3.1:** For  $n \geq 1$ ,  $D_n = n!(1 - \frac{1}{1!} + \frac{1}{2!} - \frac{1}{3!} + \dots + (-1)^n \frac{1}{n!})$

**Pf:** Use the inclusion and exclusion principle: If  $A_i \subset S$ ,  $\overline{\cup A_i} = |S| - \sum_{j=1}^n |A_j| + \sum_{i,j} |A_i \cap A_j| - \dots + (-1)^n |A_1 \cap A_2 \cap \dots \cap A_n|$ .

Choose  $S$ : What can we count which contains the set of derangements?

*Removing (all) restrictions to get something we can count*

Let  $S$  = the set of permutations of  $\{1, \dots, n\}$ . Then  $|S| = n!$ .

Choose  $A_j$  such that the set of derangements =  $\overline{\cup A_j}$ .

Let  $A_j$  = set of permutations such that  $j$  is in the  $j$ th spot.  
 $|A_j| = (n-1)!$  since there is only one choice for the  $j$ th spot (namely  $j$ ), leaving  $n-1$  terms to permute in the remaining  $n-1$  places.

$|A_i \cap A_j| = (n-2)!$  since there is only one choice for the  $i$ th spot (namely  $i$ ) and only one choice for the  $j$ th spot (namely  $j$ ), leaving  $n-2$  terms to permute in the remaining  $n-2$  places.

Similarly,  $|A_{i_1} \cap A_{i_2} \cap \dots \cap A_{i_k}| = (n-k)!$ .

Thus  $D_n = n! - \sum_{j=1}^n (n-1)! + \sum_{i,j} (n-2)! - \dots + (-1)^n (n-n)!$

$$= \binom{n}{0} n! - \binom{n}{1} (n-1)! + \binom{n}{2} (n-2)! - \dots + \binom{n}{n} (-1)^n 0!$$

$$= n! - \frac{n!}{1!} + \frac{n!}{2!} - \dots + (-1)^n \frac{n!}{n!} = n!(1 - \frac{1}{1!} + \frac{1}{2!} - \dots + (-1)^n \frac{1}{n!})$$

Recall  $\binom{n}{k}$  = number of ways to choose  $k$   $A_i$ 's.

*At least k rooks NOT on diagonal*

**Sidenote:** Finding the number of derangements is often called the hat check problem, because in the old days it was sometimes stated in the following terms: If  $n$  men check their hats, what is the probability that the hats are returned so that no one received their own hat.

Recall: If  $E \subset S$ , then the probability of  $E = P(E) = \frac{|E|}{|S|}$

$S$  = sample space,  $E$  = events.

Note: we assume each outcome is equally likely.

Suppose 4 customers at a restaurant order 4 meals. What is the probability that a waiter delivers these 4 orders to the 4 customers so that no customer receives what they ordered?

Answer:  $\frac{D_4}{4!} = 1 - 1 + \frac{1}{2} - \frac{1}{6} + \frac{1}{24} = \frac{9}{24} = 0.375$

The probability that a permutation of  $\{1, \dots, n\}$  is a derangement =  $\frac{D_n}{n!} = 1 - \frac{1}{1!} + \frac{1}{2!} - \dots + (-1)^n \frac{1}{n!}$

Recall Taylor's expansion from Calculus I,

$$f(x) = \sum_{j=0}^{\infty} \frac{f^{(j)}(a)}{j!} (x-a)^j \text{ (under appropriate hypothesis).}$$

Thus  $e^{-1} = \sum_{j=0}^{\infty} \frac{(-1)^j}{j!}$  (let  $f(x) = e^x, x = -1, a = 0$ ).

Thus  $e^{-1}$  is a good approximation for the probability of a derangement for  $n$  (slightly) large.

Thus the probability of a derangement is about the same when  $n = 5$  as it is for  $n = 50000000000$ .

$$\frac{D_5}{5!} = 0.36, \frac{D_6}{6!} = 0.36805, \frac{D_7}{7!} = 0.36785714285, e^{-1} = 0.36787944117...$$

*Does NOT depend on n*

We can derive a recursive formula for  $D_n$  (we will look at many recursive formulas in chapter 7).

Lemma A:  $D_n = (n-1)(D_{n-2} + D_{n-1})$  for  $n \geq 3$ .

Note the above formula is a recursive formula as we can determine  $D_n$  by calculating  $D_k$  for  $k < n$ .

Note  $D_1 = 0$ ,  $D_2 = 1$  (as 2 1 is the only derangement of  $\{1, 2\}$ ).

Thus  $D_3 = 2(0+1) = 2$ ,  $D_4 = 3(1+2) = 9$ ,  $D_5 = 4(2+9) = 44$ , etc.

$$\begin{aligned}
 & (3-1)(0+1) \quad (4-1)(1+2) \quad D_6 = 5(9+44) \\
 & = 2 \quad = 9 \quad = 5(53) \\
 & = 265 \\
 & = D_6
 \end{aligned}$$

Combinatorial proof of lemma A:

Let  $\mathcal{D}_n$  = the set of derangements of  $\{1, \dots, n\}$ .

$D_n$  = the number of derangements of  $\{1, \dots, n\} = |\mathcal{D}_n|$ .

We need to show that  $D_n$  is a product of  $n-1$  and  $D_{n-2} + D_{n-1}$ . If we can partition  $\mathcal{D}_n$  into  $n-1$  subsets where each subset has  $D_{n-2} + D_{n-1}$  elements, we can use the multiplication principle to show  $D_n = (n-1)(D_{n-2} + D_{n-1})$ .

We also want to relate  $D_n$  to  $D_{n-1}$  = the number of derangements of  $\{1, \dots, n-1\}$  (and  $D_{n-2}$ ).

Let's focus on one of the positions of a derangement. The last ( $n$ th) position of our derangement can be anything except  $n$ . Thus there are  $n-1$  choices for the last ( $n$ th) position. Note the factor  $n-1$  appears in our formula.

Let  $\mathcal{R}_k$  = the set of derangements of  $\{1, \dots, n\}$  where  $k$  is in the  $n$ th position for  $k = 1, \dots, n-1$ .

Then  $\mathcal{D}_n = \cup_{j=0}^{n-1} \mathcal{R}_n$

Let  $r_k = |\mathcal{R}_k|$  the number of derangements such that  $k$  is in the  $n$ th position.

Note that  $r_1 = r_2 = \dots = r_{n-1}$  (while  $r_n = 0$ ).

Then  $D_n = r_1 + \dots + r_{n-1} = r_{n-1} + \dots + r_{n-1} = (n-1)r_{n-1}$ .

Thus we have (hopefully) simplified our problem to showing that  $D_{n-2} + D_{n-1} = r_{n-1}$  = the number of derangements such that  $n-1$  is in the  $n$ th position.

We need to partition the permutations in  $\mathcal{R}_{n-1}$  into two sets, one with  $D_{n-2}$  elements and the other with  $D_{n-1}$  elements.

We can easily take care of  $D_{n-2}$ . The numbers  $n-1$  and  $n$  do not appear in any derangement of  $\{1, \dots, n-2\}$ . In  $\mathcal{R}_{n-1}$ ,  $n-1$  appears in the last position. We can take a look at the derangements in  $\mathcal{R}_{n-1}$ , such that  $n$  appears in the  $(n-1)$ st position. If we remove the  $n$ th and  $(n-1)$ st entries, we obtain a derangement in  $\mathcal{D}_{n-2}$ .

Ex: for  $n = 5$ ,  $23154 \in \mathcal{R}_{n-1} \rightarrow 231 \in \mathcal{D}_{n-2}$ .

Thus  $D_{n-2}$  = the number of derangements of  $\mathcal{R}_{n-1}$  such that  $n$  is in the  $(n-1)$ st position (and by definition of  $\mathcal{R}_{n-1}$ ,  $n-1$  is in the  $n$ th position).

We can now look at the remaining derangements in  $\mathcal{R}_{n-1}$  where  $n$  is not in the  $(n-1)$ st position.