

Scalar Line Integrals:

Let  $\mathbf{x} : [a, b] \rightarrow \mathbf{R}^n$  be a  $C^1$  path.  $f : \mathbf{R}^n \rightarrow R$ , a scalar field.

$$\begin{aligned} \Delta s_k &= \text{length of } k\text{th segment of path} \\ &= \int_{t_{k-1}}^{t_k} \|\mathbf{x}'(t)\| dt = \|\mathbf{x}'(t_k^{**})\| (t_k - t_{k-1}) = \|\mathbf{x}'(t_k^{**})\| \Delta t_k \\ &\quad \text{for some } t_k^{**} \in [t_{k-1}, t_k] \end{aligned}$$

$$\int_{\mathbf{x}} f \, ds \sim \sum_{i=1}^n f(\mathbf{x}(t_k^*)) \Delta s_k = \sum_{i=1}^n f(\mathbf{x}(t_k^*)) \|\mathbf{x}'(t_k^{**})\| \Delta t_k$$

$$\text{Thus } \int_{\mathbf{x}} f \, ds = \int_a^b f(\mathbf{x}(t)) \|\mathbf{x}'(t)\| dt$$

---

Vector Line integrals:

Let  $\mathbf{x} : [a, b] \rightarrow \mathbf{R}^n$  be a  $C^1$  path.  $F : \mathbf{R}^n \rightarrow R^n$ , a vector field.

$$\mathbf{x}'(t_k^*) \sim \frac{\Delta \mathbf{x}_k}{\Delta t_k}$$

$$\int_{\mathbf{x}} F \cdot ds \sim \sum_{i=1}^n F(\mathbf{x}(t_k^*)) \cdot \Delta \mathbf{x}_k = \sum_{i=1}^n F(\mathbf{x}(t_k^*)) \cdot \mathbf{x}'(t_k^*) \Delta t_k$$

$$\text{Thus } \int_{\mathbf{x}} F \cdot ds = \int_a^b F(\mathbf{x}(t)) \cdot \mathbf{x}'(t) dt$$

Other Formulations of Vector Line integrals:

The tangent vector to  $\mathbf{x}$  at  $t$  is  $T(t) = \frac{\mathbf{x}'(t)}{\|\mathbf{x}'(t)\|}$

$$\begin{aligned} \int_{\mathbf{x}} F \cdot ds &= \int_a^b F(\mathbf{x}(t)) \cdot \mathbf{x}'(t) dt = \int_a^b F(\mathbf{x}(t)) \cdot \mathbf{x}'(t) \frac{\|\mathbf{x}'(t)\|}{\|\mathbf{x}'(t)\|} dt = \\ &= \int_a^b F(\mathbf{x}(t)) \cdot T(t) \|\mathbf{x}'(t)\| dt = \int_{\mathbf{x}} F(\mathbf{x}(t)) \cdot T(t) ds \end{aligned}$$

Note  $\int_a^b F(\mathbf{x}(t)) \cdot T(t) ds$  is a scalar line integral of the scalar field  $F \cdot T : \mathbf{R}^n \rightarrow \mathbf{R}$  over the path  $\mathbf{x}$ .

Note:  $F \cdot T$  is the tangential component of  $F$  along the path  $\mathbf{x}$ .

---

Another notation (differential form):

For simplicity, we will work in  $\mathbf{R}^2$ , but the following generalizes to any dimension.

Let  $\mathbf{x}(t) = (x(t), y(t))$ . Let  $F(x, y) = (M(x, y), N(x, y))$

$x = x(t), y = y(t)$ . Also  $x'(t) = \frac{dx}{dt}, y'(t) = \frac{dy}{dt}$

$$\begin{aligned} \int_{\mathbf{x}} F \cdot ds &= \int_a^b F(\mathbf{x}(t)) \cdot \mathbf{x}'(t) dt \\ &= \int_a^b (M(x, y), N(x, y)) \cdot (x'(t), y'(t)) dt \\ &= \int_a^b (M(x, y), N(x, y)) \cdot (x'(t) dt, y'(t) dt) \\ &= \int_{\mathbf{x}} (M(x, y), N(x, y)) \cdot (dx, dy) \\ &= \int_{\mathbf{x}} M(x, y) dx + N(x, y) dy \end{aligned}$$

Definitions:

A *curve* is the image of piecewise  $C^1$  path  $\mathbf{x} : [a, b] \rightarrow \mathbf{R}^n$ .

A curve is *simple* if it has no self-intersections; that is,  $\mathbf{x}$  is 1:1 on the open interval  $(a, b)$

A path is *closed* if  $\mathbf{x}(a) = \mathbf{x}(b)$

A curve is *closed* if there exists a parametrization of the curve such that  $\mathbf{x}(a) = \mathbf{x}(b)$

$\int_{\mathbf{x}} F \cdot ds$  is called the *circulation* of  $f$  along  $\mathbf{x}$  if  $\mathbf{x}$  is a closed path.

A *parametrization* of a curve  $C$  is a path whose image is  $C$ . Normally we will require a parametrization of a curve to be 1:1 where possible.

---

A piecewise  $C^1$  path  $\mathbf{y} : [c, d] \rightarrow \mathbf{R}^n$  is a *reparametrization* of a piecewise  $C^1$  path  $\mathbf{x} : [a, b] \rightarrow \mathbf{R}^n$  if there exists a bijective  $C^1$  function  $u : [c, d] \rightarrow [a, b]$  where the inverse of  $u$  is also  $C^1$  and  $\mathbf{y} = \mathbf{x} \circ u$  (i.e.,  $\mathbf{y}(t) = \mathbf{x}(u(t))$ ).

Note that either

1.)  $u(a) = c$  and  $u(b) = d$ . In this case, we say that  $\mathbf{y}$  (and  $u$  are orientation-preserving OR

2.)  $u(a) = d$  and  $u(b) = c$ . In this case, we say that  $\mathbf{y}$  (and  $u$  are orientation-reversing.

---

Given piecewise  $C^1$  path,  $\mathbf{x} : [a, b] \rightarrow \mathbf{R}^n$ , the *opposite path* is  $\mathbf{x}_{opp} : [a, b] \rightarrow \mathbf{R}^n$   $\mathbf{x}_{opp} = \mathbf{x}(a + b - t)$

That is  $\mathbf{x}_{opp}$  is an orientation-reversing reparametrization of  $\mathbf{x}$  where  $u : [a, b] \rightarrow [a, b]$ ,  $u(t) = a + b - t$ .

---

Thm: Let  $\mathbf{x} : [a, b] \rightarrow \mathbf{R}^n$  be a piecewise  $C^1$  path and let  $\mathbf{y} : [c, d] \rightarrow \mathbf{R}^n$  be a reparametrization of  $\mathbf{x}$ . Then

if  $f : \mathbf{R}^n \rightarrow \mathbf{R}$  is continuous, then  $\int_{\mathbf{y}} f \, ds = \int_{\mathbf{x}} f \, ds$

if  $F : \mathbf{R}^n \rightarrow \mathbf{R}^n$  is continuous, then

$$\int_{\mathbf{y}} F \cdot ds = \int_{\mathbf{x}} F \cdot ds \text{ if } \mathbf{y} \text{ is orientation-preserving.}$$

$$\int_{\mathbf{y}} F \cdot ds = - \int_{\mathbf{x}} F \cdot ds \text{ if } \mathbf{y} \text{ is orientation-reversing.}$$