

Choose 10 of the following 11 problems. If you choose fewer than 10, the points for the remaining problems will be appropriately averaged. 3 of the problems have 1 point extra credit available (indicated by [1-EC]). You can do the extra credit even if you do not choose these problems.

[3] 1a.) Without using the LaPlace transform, solve the following initial value problem:

$$y'' + 2y' + y = 0, \quad y(0) = 0, \quad y'(0) = 4$$

Answer 1a.) \_\_\_\_\_

[4] 1b.) Without using the LaPlace transform, find the general solution to the differential equation  $y'' + 2y' + y = e^{-t}$

Answer 1b.) \_\_\_\_\_

[3] 1c.) Without using the LaPlace transform, solve the following initial value problem:

$$y'' + 2y' + y = e^{-t}, \quad y(0) = 0, \quad y'(0) = 4$$

Answer 1c.) \_\_\_\_\_

[10] 2.) Use the LaPlace transform to solve the following initial value problem:

$$y'' + 2y' + y = e^{-t}, \quad y(0) = 0, \quad y'(0) = 4$$

Answer 2.) \_\_\_\_\_

[1-EC] 2a.) How can you use the LaPlace transform to find the general solution to a differential equation.

[10] 3.) Find the inverse Laplace transform of  $e^{-3s} \frac{s}{s^2+8s+18}$

Answer 3.) \_\_\_\_\_

[10] 4.) Use the definition and not the table to find the Laplace transform of  $f(t) = 3t^2$ .  
CLEARLY indicate when you are taking a limit.

Answer 4.) \_\_\_\_\_

[5] 5a.) Find the inverse Laplace transform  $\frac{1}{(s-3)(s^2-6s+10)}$  by using the convolution integral. Leave your answer in terms of a convolution integral:

Answer 5a.) \_\_\_\_\_

[5] 5b.) Evaluate the convolution integral obtained in 3a.

Answer 5b.) \_\_\_\_\_

[10] 6.) Solve the following initial value problem:  $\frac{y'}{t^2} = -3y + 1$ ,  $y(0) = 0$

Answer 6.) \_\_\_\_\_

[10] 7.) Draw the direction field for the differential equation  $y' = (y - 2)(y + 2)$ . Based on the direction field, determine the behavior of  $y$  as  $t \rightarrow \infty$ . If this behavior depends on the initial value of  $y$  at  $t = 0$ , describe this dependency.

[5] 8a.) Transform  $x'' - 2x' - 3x = e^t$  into a system of first order differential equations.

Answer 8a.) \_\_\_\_\_

[5] 8b.) Use Euler's formula to write  $e^{4-2i}$  in the form of  $a + ib$ .

Answer 8b.) \_\_\_\_\_

9.) Circle T for true and F for false.

[2] 9a.) Given an initial value, there always exists a unique solution to any second order differential equation.

T F

[2] 9b.) Given an initial value, there always exists at least one solution to any second order differential equation.

T F

[2] 9c.) If there is no damping and no external force, then the general solution to a mechanical vibration problem with constant spring force must be of the form  $c_1 \cos(\omega_0 t) + c_2 \sin(\omega_0 t)$

T F

[1] 9d.)  $y = c_1 e^{3t} + c_2 t e^{3t} + 4 \cos t$  is a possible general solution to a mechanical vibration problem with damping.

T F

[4 or 1-EC] 9e.) Explain your answer to problem 9d.

[10] 10a.) A mass weighing 2 kg stretches a spring .1m. If the mass is pulled down an additional .2m and released, and if there is no damping, determine the position of the mass at any time  $t$

Answer 10a.) \_\_\_\_\_

[1-EC] 10b.) Do the initial conditions affect the long-term behavior of the motion of the mass?

[10] 11.) Suppose that a sum  $S_0$  is invested at an annual rate of return  $r$  compounded continuously. Find the time  $T$  required for the original sum to double in value as a function of  $r$ . Assume that the rate of change of the value of the investment is equal to the interest rate  $r$  times the current value of the investment  $S(t)$ .

Answer 11.) \_\_\_\_\_