

Suppose  $A \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \lambda_1 \begin{bmatrix} v_1 \\ v_2 \end{bmatrix}$  and  $A \begin{bmatrix} w_1 \\ w_2 \end{bmatrix} = \lambda_2 \begin{bmatrix} w_1 \\ w_2 \end{bmatrix}$

Claim:

If  $A = A^T$  and  $\lambda_1 \neq \lambda_2$ , then  $\begin{bmatrix} v_1 \\ v_2 \end{bmatrix}$  is perpendicular to  $\begin{bmatrix} w_1 \\ w_2 \end{bmatrix}$

I.e., If eigenvectors come from different eigenspaces, then the eigenvectors are orthogonal WHEN  $A = A^T$ .

$$\begin{aligned} \text{Pf of claim: } \lambda_1(v_1, v_2) \cdot (w_1, w_2) &= \lambda_1[v_1, v_2] \begin{bmatrix} w_1 \\ w_2 \end{bmatrix} \\ &= (\lambda_1 \begin{bmatrix} v_1 \\ v_2 \end{bmatrix})^T \begin{bmatrix} w_1 \\ w_2 \end{bmatrix} = (A \begin{bmatrix} v_1 \\ v_2 \end{bmatrix})^T \begin{bmatrix} w_1 \\ w_2 \end{bmatrix} \\ &= \begin{bmatrix} v_1 \\ v_2 \end{bmatrix}^T A^T \begin{bmatrix} w_1 \\ w_2 \end{bmatrix} = [v_1, v_2] A \begin{bmatrix} w_1 \\ w_2 \end{bmatrix} \\ &= [v_1, v_2] \lambda_2 \begin{bmatrix} w_1 \\ w_2 \end{bmatrix} = \lambda_2[v_1, v_2] \begin{bmatrix} w_1 \\ w_2 \end{bmatrix} \\ &= \lambda_2(v_1, v_2) \cdot (w_1, w_2) \end{aligned}$$

$$\lambda_1(v_1, v_2) \cdot (w_1, w_2) = \lambda_2(v_1, v_2) \cdot (w_1, w_2)$$

$$\text{implies } \lambda_1(v_1, v_2) \cdot (w_1, w_2) - \lambda_2(v_1, v_2) \cdot (w_1, w_2) = 0.$$

$$\text{Thus } (\lambda_1 - \lambda_2)(v_1, v_2) \cdot (w_1, w_2) = 0$$

$$\lambda_1 \neq \lambda_2 \text{ implies } (v_1, v_2) \cdot (w_1, w_2) = 0$$

Thus these eigenvectors are orthogonal.

## 7.1: Orthogonal Diagonalization

Equivalent Questions:

- Given an  $n \times n$  matrix, does there exist an orthonormal basis for  $R^n$  consisting of eigenvectors of  $A$ ?
- Given an  $n \times n$  matrix, does there exist an orthonormal matrix  $P$  such that  $P^{-1}AP = P^TAP$  is a diagonal matrix?
- Is  $A$  symmetric?

Defn: A matrix is **symmetric** if  $A = A^T$ .

Recall An invertible matrix  $P$  is **orthogonal** if  $P^{-1} = P^T$

Defn: A matrix  $A$  is **orthogonally diagonalizable** if there exists an orthogonal matrix  $P$  such that  $P^{-1}AP = D$  where  $D$  is a diagonal matrix.

Thm: If  $A$  is an  $n \times n$  matrix, then the following are equivalent:

- a.)  $A$  is orthogonally diagonalizable.
- b.) There exists an orthonormal basis for  $R^n$  consisting of eigenvectors of  $A$ .
- c.)  $A$  is symmetric.

Thm: If  $A$  is a symmetric matrix, then:

- a.) The eigenvalues of  $A$  are all real numbers.
- b.) Eigenvectors from different eigenspaces are orthogonal.
- c.) Geometric multiplicity of an eigenvalue = its algebraic multiplicity

Note if  $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$  are linearly independent:

(1.) You can use the Gram-Schmidt algorithm to find an orthogonal basis for  $\text{span}\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ .

(2.) You can normalize these orthogonal vectors to create an orthonormal basis for  $\text{span}\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ .

(3.) These basis vectors are not normally eigenvectors of  $A = [\mathbf{v}_1 \dots \mathbf{v}_n]$  even if  $A$  is symmetric (note that there are an infinite number of orthogonal basis for  $\text{span}\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$  even if  $n = 2$  and  $\text{span}\{\mathbf{v}_1, \mathbf{v}_2\}$  is just a 2-dimensional plane)

---

Note if  $A$  is a  $n \times n$  square matrix that is diagonalizable, then you can find  $n$  linearly independent eigenvectors of  $A$ .

Each eigenvector is in  $\text{col}(A)$ : If  $\mathbf{v}$  is an eigenvector of  $A$  with eigenvalue  $\lambda$ , then  $A\mathbf{v} = \lambda\mathbf{v}$ . Thus  $\frac{1}{\lambda}A\mathbf{v} = \mathbf{v}$ .

Hence  $A(\frac{1}{\lambda}\mathbf{v}) = \mathbf{v}$ . Thus  $\mathbf{v}$  is in  $\text{col}(A)$ .

Thus  $\text{col}(A)$  is an  $n$ -dimensional subspace of  $R^n$ . That is  $\text{col}(A) = R^n$ , and you can find a basis for  $\text{col}(A) = R^n$  consisting of eigenvectors of  $A$ .

But these eigenvectors are NOT usually orthogonal UNLESS they come from different eigenspaces AND the matrix  $A$  is symmetric.

If  $A$  is NOT symmetric, then eigenvectors from different eigenspaces need NOT be orthogonal.

IF  $A$  is symmetric,

To orthogonally diagonalize a symmetric matrix  $A$ :

1.) Find the eigenvalues of  $A$ .

Solve  $\det(A - \lambda I) = 0$  for  $\lambda$ .

2.) Find a basis for each of the eigenspaces.

Solve  $(A - \lambda_j I)\mathbf{x} = 0$  for  $\mathbf{x}$ .

3.) Use the Gram-Schmidt process to find an orthonormal basis for each eigenspace.

That is for each  $\lambda_j$  use Gram-Schmidt to find an orthonormal basis for  $Nul(A - \lambda_j I)$ .

Eigenvectors from different eigenspaces will be orthogonal, so you don't need to apply Gram-Schmidt to eigenvectors from different eigenspaces

4.) Use the eigenvalues of  $A$  to construct the diagonal matrix  $D$ , and use the orthonormal basis of the corresponding eigenspaces for the corresponding columns of  $P$ .

5.) Note  $P^{-1} = P^T$  since the columns of  $P$  are orthonormal.

Example 1:

Orthogonally diagonalize  $A = \begin{bmatrix} 1 & 2 \\ 2 & 4 \end{bmatrix}$

Step 1: Find the eigenvalues of  $A$ :

$$\begin{aligned} \det(A - \lambda I) &= \begin{vmatrix} 1 - \lambda & 2 \\ 2 & 4 - \lambda \end{vmatrix} = (1 - \lambda)(4 - \lambda) - 4 \\ &= \lambda^2 - 5\lambda + 4 - 4 = \lambda^2 - 5\lambda = \lambda(\lambda - 5) = 0 \end{aligned}$$

Thus  $\lambda = 0, 5$  are are eigenvalues of  $A$ .

2.) Find a basis for each of the eigenspaces:

$$\lambda = 0 : (A - 0I) = A = \begin{bmatrix} 1 & 2 \\ 2 & 4 \end{bmatrix} \sim \begin{bmatrix} 1 & 2 \\ 0 & 0 \end{bmatrix}$$

Thus  $\begin{bmatrix} -2 \\ 1 \end{bmatrix}$  is an eigenvector of  $A$  with eigenvalue 0.

$$\lambda = 5 : (A - 5I) = \begin{bmatrix} -4 & 2 \\ 2 & -1 \end{bmatrix} \sim \begin{bmatrix} 2 & -1 \\ 0 & 0 \end{bmatrix}$$

Thus  $\begin{bmatrix} 1 \\ 2 \end{bmatrix}$  is an eigenvector of  $A$  with eigenvalue 5.

3.) Create orthonormal basis:

Since  $A$  is symmetric and the eigenvectors  $\begin{bmatrix} -2 \\ 1 \end{bmatrix}$  and  $\begin{bmatrix} 1 \\ 2 \end{bmatrix}$  come from different eigenspaces (ie their eigenvalues are different), these eigenvectors are orthogonal. Thus we only

need to normalize them:

$$\left\| \begin{bmatrix} -2 \\ 1 \end{bmatrix} \right\| = \sqrt{4+1} = \sqrt{5}$$

$$\left\| \begin{bmatrix} 1 \\ 2 \end{bmatrix} \right\| = \sqrt{1+4} = \sqrt{5}$$

Thus an orthonormal basis for  $\text{col}(A) = R^2 = \left\{ \begin{bmatrix} \frac{-2}{\sqrt{5}} \\ \frac{1}{\sqrt{5}} \end{bmatrix}, \begin{bmatrix} \frac{1}{\sqrt{5}} \\ \frac{2}{\sqrt{5}} \end{bmatrix} \right\}$  ■

4.) Construct  $D$  and  $P$

$$D = \begin{bmatrix} 0 & 0 \\ 0 & 5 \end{bmatrix}, \quad P = \begin{bmatrix} \frac{-2}{\sqrt{5}} & \frac{1}{\sqrt{5}} \\ \frac{1}{\sqrt{5}} & \frac{2}{\sqrt{5}} \end{bmatrix}$$

Make sure order of eigenvectors in  $D$  match order of eigenvalues in  $P$ .

5.)  $P$  orthonormal implies  $P^{-1} = P^T$

$$\text{Thus } P^{-1} = \begin{bmatrix} \frac{-2}{\sqrt{5}} & \frac{1}{\sqrt{5}} \\ \frac{1}{\sqrt{5}} & \frac{2}{\sqrt{5}} \end{bmatrix}$$

Note that in this example,  $P^{-1} = P$ , but that is NOT normally the case.

Thus  $A = PDP^{-1}$

$$\text{Thus } \begin{bmatrix} 1 & 2 \\ 2 & 4 \end{bmatrix} = \begin{bmatrix} \frac{-2}{\sqrt{5}} & \frac{1}{\sqrt{5}} \\ \frac{1}{\sqrt{5}} & \frac{2}{\sqrt{5}} \end{bmatrix} \begin{bmatrix} 0 & 0 \\ 0 & 5 \end{bmatrix} \begin{bmatrix} \frac{-2}{\sqrt{5}} & \frac{1}{\sqrt{5}} \\ \frac{1}{\sqrt{5}} & \frac{2}{\sqrt{5}} \end{bmatrix}$$

Example 2:  
 Orthogonally diagonalize  $A = \begin{bmatrix} 1 & -1 & 1 \\ -1 & 1 & -1 \\ 1 & -1 & 1 \end{bmatrix}$

Step 1: Find the eigenvalues of  $A$ :

$$\det(A - \lambda I) = \begin{vmatrix} 1 - \lambda & -1 & 1 \\ -1 & 1 - \lambda & -1 \\ 1 & -1 & 1 - \lambda \end{vmatrix} = \begin{vmatrix} 1 - \lambda & -1 & 1 \\ -1 & 1 - \lambda & -1 \\ 0 & -\lambda & -\lambda \end{vmatrix}$$

$$= (1 - \lambda) \begin{vmatrix} 1 - \lambda & -1 \\ -\lambda & -\lambda \end{vmatrix} - (-1) \begin{vmatrix} -1 & 1 \\ -\lambda & -\lambda \end{vmatrix} + 0 \begin{vmatrix} -1 & 1 \\ 1 - \lambda & -1 \end{vmatrix}$$

$$= (1 - \lambda)[(1 - \lambda)(-\lambda) - \lambda] + [\lambda + \lambda]$$

$$= (1 - \lambda)(-\lambda)[(1 - \lambda) + 1] + 2\lambda = (1 - \lambda)(-\lambda)(2 - \lambda) + 2\lambda$$

Note I can factor out  $-\lambda$ , leaving only a quadratic to factor:

$$= -\lambda[(1 - \lambda)(2 - \lambda) - 2]$$

$$= -\lambda[\lambda^2 - 3\lambda + 2 - 2] = -\lambda[\lambda^2 - 3\lambda] = -\lambda^2[\lambda - 3]$$

Thus there are 2 eigenvalues:

$\lambda = 0$  with algebraic multiplicity 2. Since  $A$  is symmetric, geometric multiplicity = algebraic multiplicity = 2. Thus the dimension of the eigenspace corresponding to  $\lambda = 0$  [ $= \text{Nul}(A - 0I) = \text{Nul}(A)$ ] is 2.

$\lambda = 3$  w/ algebraic multiplicity = 1 = geometric multiplicity.

Thus we can find an orthogonal basis for  $R^3$  where two of the basis vectors come from the eigenspace corresponding to eigenvalue 0 while the third comes from the eigenspace corresponding to eigenvalue 3.

2.) Find a basis for each of the eigenspaces:

$$2a.) \lambda = 0 : A - 0I = A = \begin{bmatrix} 1 & -1 & 1 \\ -1 & 1 & -1 \\ 1 & -1 & 1 \end{bmatrix} \sim \begin{bmatrix} 1 & -1 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} x_2 - x_3 \\ x_2 \\ x_3 \end{bmatrix} = x_2 \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} + x_3 \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix}$$

Thus a basis for eigenspace corresponding to eigenvalue 0 is

$$\left\{ \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix} \right\}$$

We can now use Gram-Schmidt to turn this basis into an orthogonal basis for the eigenspace corresponding to eigenvalue 0 or we can continue finding eigenvalues.

3a.) Create orthonormal basis using Gram-Schmidt for the eigenspace corresponding to eigenvalue 0:

$$\text{Let } \mathbf{v}_1 = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} \text{ and } \mathbf{v}_2 = \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix}$$

$$\text{proj}_{\mathbf{v}_1} \mathbf{v}_2 = \frac{\mathbf{v}_2 \cdot \mathbf{v}_1}{\mathbf{v}_1 \cdot \mathbf{v}_1} \mathbf{v}_1 = \frac{-1+0+0}{1+1+0} \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} = -\frac{1}{2} \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} -\frac{1}{2} \\ -\frac{1}{2} \\ 0 \end{bmatrix}$$



The vector component of  $\mathbf{v}_2$  orthogonal to  $\mathbf{v}_1$  is

$$\mathbf{v}_2 - \text{proj}_{\mathbf{v}_1} \mathbf{v}_2 = \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix} - \begin{bmatrix} -\frac{1}{2} \\ -\frac{1}{2} \\ 0 \end{bmatrix} = \begin{bmatrix} -\frac{1}{2} \\ \frac{1}{2} \\ 1 \end{bmatrix}$$

Thus an orthogonal basis for the eigenspace corresponding to eigenvalue 0 is

$$\left\{ \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} -\frac{1}{2} \\ \frac{1}{2} \\ 1 \end{bmatrix} \right\}$$

To create orthonormal basis, divide each vector by its length:

$$\left\| \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} \right\| = \sqrt{1^2 + 1^2 + 0^2} = \sqrt{2}$$

$$\left\| \begin{bmatrix} -\frac{1}{2} \\ \frac{1}{2} \\ 1 \end{bmatrix} \right\| = \sqrt{\frac{1}{4} + \frac{1}{4} + 1} = \sqrt{\frac{3}{2}}$$

Thus an orthonormal basis for the eigenspace corresponding to eigenvalue 0 is

$$\left\{ \begin{bmatrix} \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} \\ 0 \end{bmatrix}, \begin{bmatrix} -\frac{\sqrt{2}}{2\sqrt{3}} \\ \frac{\sqrt{2}}{2\sqrt{3}} \\ \sqrt{\frac{2}{3}} \end{bmatrix} \right\} = \left\{ \begin{bmatrix} \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} \\ 0 \end{bmatrix}, \begin{bmatrix} -\frac{1}{\sqrt{6}} \\ \frac{1}{\sqrt{6}} \\ \sqrt{\frac{2}{3}} \end{bmatrix} \right\}$$

2b.) Find a basis for eigenspace corresponding to  $\lambda = 3$  :

$$A-3I = \begin{bmatrix} -2 & -1 & 1 \\ -1 & -2 & -1 \\ 1 & -1 & -2 \end{bmatrix} \sim \begin{bmatrix} 1 & -1 & -2 \\ 0 & -3 & -3 \\ 0 & -3 & -3 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & -1 \\ 0 & 1 & 1 \\ 0 & 0 & 0 \end{bmatrix}$$

Thus a basis for eigenspace corresponding to eigenvalue 3 is

$$\left\{ \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \right\}$$

FYI: Alternate method to find 3rd vector: Since you have two linearly independent vectors from the eigenspace corresponding to eigenvalue 0, you only need one more vector which is orthogonal to these two to form a basis for  $R^3$ . Note since  $A$  is symmetric, any such vector will be an eigenvector of  $A$  with eigenvalue 3. Note this shortcut only works because we know what the eigenspace corresponding to eigenvalue 3 looks like: a line perpendicular to the plane representing the eigenspace corresponding to eigenvalue 0.

3b.) Create orthonormal basis for the eigenspace corresponding to eigenvalue 3:

We only need to normalize:

$$\left\| \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \right\| = \sqrt{1^2 + (-1)^2 + 1^2} = \sqrt{3}$$

Thus orthonormal basis for eigenspace corresponding to eigenvalue 3 is

$$\left\{ \begin{bmatrix} \frac{1}{\sqrt{3}} \\ -\frac{1}{\sqrt{3}} \\ \frac{1}{\sqrt{3}} \end{bmatrix} \right\}$$

4.) Construct  $D$  and  $P$

$$D = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 3 \end{bmatrix}, \quad P = \begin{bmatrix} \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{6}} & -\frac{1}{\sqrt{3}} \\ 0 & \sqrt{\frac{2}{3}} & \frac{1}{\sqrt{3}} \end{bmatrix}$$

Make sure order of eigenvectors in  $D$  match order of eigenvalues in  $P$ .

5.)  $P$  orthonormal implies  $P^{-1} = P^T$

$$\text{Thus } P^{-1} = \begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} & 0 \\ -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{6}} & \sqrt{\frac{2}{3}} \\ \frac{1}{\sqrt{3}} & -\frac{1}{\sqrt{3}} & \frac{1}{\sqrt{3}} \end{bmatrix}$$

$$\text{Thus } \begin{bmatrix} 1 & -1 & 1 \\ -1 & 1 & -1 \\ 1 & -1 & 1 \end{bmatrix} = A = PDP^{-1} =$$

$$\begin{bmatrix} \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{6}} & -\frac{1}{\sqrt{3}} \\ 0 & \sqrt{\frac{2}{3}} & \frac{1}{\sqrt{3}} \end{bmatrix} \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 3 \end{bmatrix} \begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} & 0 \\ -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{6}} & \sqrt{\frac{2}{3}} \\ \frac{1}{\sqrt{3}} & -\frac{1}{\sqrt{3}} & \frac{1}{\sqrt{3}} \end{bmatrix}$$