

# STAT:2010/4200 Statistical Methods and Computing

## Introduction to Hypothesis Testing

Lecture 14  
Mar. 11, 2020

Kate Cowles  
374 SH, 335-0727  
kate-cowles@uiowa.edu

Example:

I claim that my husband's resting pulse rate is 45 beats per minute. This is very low and would be typical of either a highly trained athlete or a sick individual.

To test my claim, you wish to measure his resting heart rate on 5 different occasions.

Here, the "population" of interest is all possible measurements of my husband's resting pulse rate. My claim may be interpreted as saying that the mean  $\mu$  of this "population" of values is 45 beats per minute.

## Introduction to Hypothesis Testing

Recall that statistical inference is using data contained in a sample to draw conclusions or make decisions about the entire population from which the sample is taken.

Two main goals of statistical inference

- estimation of unknown population parameters
- testing specific hypotheses about unknown population parameters

The purpose of hypothesis testing is to "assess the evidence provided by data about some claim concerning a population."\*

\* Moore, D.S. *The Basic Practice of Statistics*

Suppose the measurements you get are:

42    52    43    48    47

The sample mean  $\bar{x} = 46.4$ . Does this provide evidence against my claim?

We will consider this question by asking what would happen if my claim were true and we repeated the sample of 5 measurements many times.

Suppose first that we knew that the standard deviation of measurements of my husband's resting heart rate was  $\sigma = 4$  beats per minute.

- If the claim that  $\mu = 45$  is true, the sampling distribution of  $\bar{x}$  from 5 measurements is normal with mean  $\mu = 45$  and standard deviation

$$\frac{\sigma}{\sqrt{n}} = \frac{4}{\sqrt{5}} = 1.79$$

- We can judge whether any observed  $\bar{x}$  is surprising by finding it on this distribution.

The *alternative hypothesis* is the claim *for* which we are trying to find evidence.

- symbolized  $H_a$

In the example about my husband's heart rate, your alternative hypothesis probably was

$$H_a : \mu > 45$$

The *p-value* of the test is the probability, computed assuming that  $H_0$  is true, that the observed outcome would take a value as extreme as or more extreme than, what we actually observed.

- Small p-values are evidence against the null hypothesis.

## Terminology of hypothesis tests

The *null hypothesis* is the statement being tested.

- The test is intended to assess the strength of evidence *against* the null hypothesis.
- Usually is a statement of "no effect," "no difference," "nothing going on."
- The null hypothesis is commonly symbolized as  $H_0$ .
- $H_0$  is a statement about an unknown population parameter.
- Example:

$$H_0 : \mu = 45$$

The result of a hypothesis test is a decision. The possible outcomes are called

- Rejecting the null hypothesis
- Not rejecting the null hypothesis

*Before* we carry out the test, we must decide how strong we will require the evidence to be in order for us to reject  $H_0$ . We specify this in terms of a *significance level*.

- The significance level is how small we will require the p-value to be in order to reject  $H_0$ .
- symbol is  $\alpha$
- conventional choices are  $\alpha = .05$  and  $\alpha = .01$

Example: my husband's resting heart rate

We will choose  $\alpha = .05$  as the significance level at which to carry out the test.

To find the p-value of our results, we will standardize  $\bar{x}$  so we can use the normal table.

- Remember: the p-value is computed assuming  $H_0$  is true, so the value of  $\mu$  to use is the value stated in  $H_0$ .

$$\begin{aligned} z &= \frac{\bar{x} - \mu}{\sigma/\sqrt{n}} \\ &= \frac{46.4 - 45}{1.79} \\ &= 0.78 \end{aligned}$$

## One-sided and two-sided tests of hypotheses

The hypothesis test we just conducted was *one-sided* test. We were interested only in showing that the value of the unknown parameter differed from that given in  $H_0$  in one direction.

$$\begin{aligned} H_0 &: \mu = 45 \\ H_a &: \mu > 45 \end{aligned}$$

We might also have stated the hypotheses this way:

$$\begin{aligned} H_0 &: \mu \leq 45 \\ H_a &: \mu > 45 \end{aligned}$$

According to Table A, the probability of a value this large or larger is 0.218. We would say that for this test result

$$p = 0.218$$

Since this is *larger* than  $\alpha = .05$ , we cannot reject the null hypothesis. That is, we have decided that the evidence was not sufficient to reject my claim!

In specifying null and alternative hypotheses:

- There must be no overlap in the range of values included in the two hypotheses.
- All possible values of the unknown population parameter must be covered in one or the other of the two hypotheses.

## Two-sided hypothesis tests

Example: We wish to compare fasting serum cholesterol levels in persons over 21 living in a group of islands in the South Pacific with typical levels found in the U.S.

We know that levels in adults over 21 in the US are approximately normally distributed with

- mean 190 mg/dl
- standard deviation 40 mg/dl.

We have no idea what the relative levels of serum cholesterol are on the islands as compared with the U.S.

The hypotheses for our *two-sided* test are:

$$H_0 : \mu = 190$$

$$H_a : \mu \neq 190$$

Before we look at our data, we will decide on the *significance level*  $\alpha$  for our test. Let us choose  $\alpha = .05$ .

We then perform blood tests on 100 adults from the islands and find that the sample mean level  $\bar{x} = 181.5$  mg/dl.

To carry out our hypothesis test, we note that, if  $H_0$  is true, the sampling distribution of  $\bar{x}$  is normal with

$$\mu = 190$$

$$\sigma_{\bar{x}} = \frac{40}{\sqrt{100}} = 4$$

We will assume that the levels on the islands are normally distributed with

- unknown mean  $\mu$
- known standard deviation 40 mg/dl

We will standardize the value of  $\bar{x}$  that we observed to find out how likely we would have been to get a value as extreme as what we got, or more extreme, if  $H_0$  were true.

$$z = \frac{\bar{x} - \mu_0}{\sigma/\sqrt{n}}$$

$$= \frac{181.5 - 190}{4}$$

$$= -2.125$$

We must find out what area under the standard normal curve lies

- to the left of -2.125
- *and* to the right of 2.125

The answer is  $.017 + .017 = .034$ .

This is the  $p$  – *value* for the test. Since  $p < .05$  we reject the null hypothesis and conclude that serum cholesterol levels are different among adult residents of the Pacific Islands than among adults in the U.S.

$$= \frac{181.5 - 190}{3.81}$$

$$= -2.231$$

## One sample t-tests

If we don't know the population standard deviation, then we

- estimate it with the sample standard deviation  $s$
- compute a  $t$  statistic rather than a  $z$  statistic
- compare to a  $t$  distribution with the appropriate degrees of freedom

Example: If we do *not* assume that we know  $\sigma$  for serum cholesterol levels among residents of the Pacific Islands.

From the sample of 100 adults, we compute

$$s = 38.1 \text{ mg/dl}$$

We then compute

$$t = \frac{\bar{x} - \mu_0}{s/\sqrt{n}}$$

We try to use Table C to find the area to the left of -2.231 and to the right of 2.231 under a  $t$  curve with 99 degrees of freedom.

The closest we can come is that under a  $t$  curve with 100 degrees of freedom, the area in one tail would be between .01 and .02.

Thus we conclude that the p-value is somewhere between .02 and .04.

SAS can do a much better job for us! It would provide a p-value of .0279.

Thus, if we had chosen  $\alpha = .05$ , we would reject the null hypothesis.

## Types of error in hypothesis testing

$$H_0 : \mu = \mu_0$$

$$H_a : \mu \neq \mu_0$$

True state of the world

	$H_0$ is false	$H_0$ is true
Reject $H_0$	Correct!	Type I error
Do not reject $H_0$	Type II error	Correct!

$$\alpha = P(\text{reject } H_0 \mid H_0 \text{ is true})$$

$$\beta = P(\text{fail to reject } H_0 \mid H_0 \text{ is false})$$

or, put another way

$\alpha$  = probability of making Type I error

$\beta$  = probability of making Type II error

power  $(1 - \beta)$  = probability of correctly rejecting  $H_0$  when it is false; depends on our definition of  $H_a$

Return to the example of my husband's resting heart rate.

- What value of  $\bar{x}$  would have been required in order to reject

$$H_0 : \mu = 45$$

in favor of

$$H_a : \mu > 45$$

if  $\alpha = .05$ ?

For a standard normal,  $z = 1.645$  cuts off the upper .05 area.

The corresponding value for the sampling distribution of  $\bar{x}$  if  $H_0$  is true is

$$\bar{x} = \mu + z\sigma$$

$$= 45 + 1.645(1.79)$$

$$= 47.9$$