

# Density Estimation and Smoothing

## Density Estimation

- Suppose we have a random sample  $X_1, \dots, X_n$  from a population with density  $f$ .
- Nonparametric density estimation is useful if we
  - want to explore the data without a specific parametric model
  - want to assess the fit of a parametric model
  - want a compromise between a parametric and a fully non-parametric approach

- A simple method for estimating  $f$  at a point  $x$ :

$$\hat{f}_n(x) = \frac{\text{no. of } X_i \text{ in } [x-h, x+h]}{2hn}$$

for some small value of  $h$

- This estimator has bias

$$\text{Bias}(\hat{f}_n(x)) = \frac{1}{2h}p_h(x) - f(x)$$

and variance

$$\text{Var}(\hat{f}_n(x)) = \frac{p_h(x)(1 - p_h(x))}{4h^2n}$$

with

$$p_h(x) = \int_{x-h}^{x+h} f(u)du$$

- If  $f$  is continuous at  $x$  and  $f(x) > 0$ , then as  $h \rightarrow 0$ 
  - the bias tends to zero;
  - the variance tends to infinity.
- Choosing a good value of  $h$  involves a *variance-bias tradeoff*.

## Kernel Density Estimation

- The estimator  $\hat{f}_n(x)$  can be written as

$$\hat{f}_n(x) = \frac{1}{nh} \sum_{i=1}^n K\left(\frac{x - x_i}{h}\right)$$

with

$$K(u) = \begin{cases} 1/2 & \text{if } |u| < 1 \\ 0 & \text{otherwise} \end{cases}$$

- Other *kernel functions*  $K$  can be used; usually
  - $K$  is a density function
  - $K$  has mean zero
  - $K$  has positive, finite variance  $\sigma_K^2$

Often  $K$  is symmetric.

- Common choices of  $K$ :

$K(u)$	Range	Name
$1/2$	$ u  < 1$	Uniform, Boxcar
$\frac{1}{\sqrt{2\pi}} e^{-u^2/2}$		Gaussian
$1 -  u $	$ u  < 1$	Triangular
$\frac{3}{4}(1 - u^2)$	$ u  < 1$	Epanechnikov
$\frac{15}{16}(1 - u^2)^2$	$ u  < 1$	Biweight

## Mean Square Error for Kernel Density Estimators

- The bias and variance of a kernel density estimator are of the form

$$\begin{aligned}\text{Bias}(\hat{f}_n(x)) &= \frac{h^2 \sigma_K^2 f''(x)}{2} + O(h^4) \\ \text{Var}(\hat{f}_n(x)) &= \frac{f(x)R(K)}{nh} + o\left(\frac{1}{nh}\right)\end{aligned}$$

with

$$R(g) = \int g(x)^2 dx$$

if  $h \rightarrow 0$  and  $nh \rightarrow \infty$  and  $f$  is reasonable.

- The pointwise asymptotic mean square error is

$$\text{AMSE}(\hat{f}_n(x)) = \frac{f(x)R(K)}{nh} + \frac{h^4 \sigma_K^4 f''(x)^2}{4}$$

and the asymptotic mean integrated square error is

$$\text{AMISE}(\hat{f}_n) = \frac{R(K)}{nh} + \frac{h^4 \sigma_K^4 R(f'')}{4}$$

- The resulting asymptotically optimal bandwidths  $h$  are

$$\begin{aligned}h_0(x) &= \left( \frac{f(x)R(K)}{\sigma_K^4 f''(x)^2} \right)^{1/5} n^{-1/5} \\ h_0 &= \left( \frac{R(K)}{\sigma_K^4 R(f'')} \right)^{1/5} n^{-1/5}\end{aligned}$$

with optimal AMSE and AMISE

$$\begin{aligned}\text{AMSE}_0(\hat{f}_n(x)) &= \frac{5}{4} (\sigma_K f(x) R(K))^{4/5} f''(x)^{2/5} n^{-4/5} \\ \text{AMISE}_0(\hat{f}_n) &= \frac{5}{4} (\sigma_K R(K))^{4/5} R(f'')^{1/5} n^{-4/5}\end{aligned}$$

## Choosing a Bandwidth

- One way to choose a bandwidth is to target a particular family, such as a Gaussian  $f$ :

- The optimal bandwidth for minimizing AMISE when  $f$  is Gaussian and  $K$  is Gaussian

$$h_0 = 1.059\sigma n^{-1/5}$$

- $\sigma$  can be estimated using  $S$  or the interquartile range
- The default for `density` in R is

$$0.9 \times \min(S, \text{IQR}/1.34) n^{-1/5}$$

based on a suggestion of Silverman (1986, pp 45–47).

- This can often serve as a reasonable starting point.
- It does not adapt to information in the data that suggests departures from normality.
- So-called *plug-in* methods estimate  $R(f'')$  to obtain

$$\hat{h} = \left( \frac{R(K)}{\sigma_K^4 \widehat{R(f'')}} \right)^{1/5} n^{-1/5}$$

- The Sheather-Jones method uses a different bandwidth (and kernel?) to estimate  $\hat{f}$  and then estimates  $R(f'')$  by  $R(\hat{f}'')$ .
- Specifying `bw="SJ"` in R's `density` uses the Sheather-Jones method. There are two variants:
  - `SJ-dpi`: direct plug-in
  - `SJ-ste`: solve the equation


The default for `bw="SJ"` is `ste`.

- Other approaches based on leave-one-out cross-validation are available.
- Many of these are available as options in R's `density` and/or other density estimation functions available in R packages.
- Variable bandwidth approaches can be based on pilot estimates of the density produced with simpler fixed bandwidth rules.

## Example: Durations of Eruptions of Old Faithful

- Based on an example in Venables and Ripley (2002).
- Durations, in minutes, of 299 consecutive eruptions of Old Faithful were recorded.
- The data are available as data set `geyser` in package `MASS`.
- Some density estimates are produced by

```
library(MASS)
data(geyser)
truehist(geyser$duration, nbin=25, col="lightgrey")
lines(density(geyser$duration))
lines(density(geyser$duration, bw="SJ"), col="red")
lines(density(geyser$duration, bw="SJ-dpi"), col="blue")
```



- Animation can be a useful way of understanding the effect of smoothing parameter choice. See files `tkdens.R`, `shinydens.R`, and `geyser.R` in

`http://www.stat.uiowa.edu/~luke/classes/STAT7400-2020/examples/`

Also

`http:  
//www.stat.uiowa.edu/~luke/classes/STAT7400-2020/examples/smoothex.Rmd`

## Issues and Notes

- Kernel methods do not work well at boundaries of bounded regions.
- Transforming to unbounded regions is often a good alternative.
- Variability can be assessed by asymptotic methods or by bootstrapping.
- A crude MCMC bootstrap animation:

```
g <- geyser$duration
for (i in 1:1000) {
  g[sample(299,1)] <- geyser$duration[sample(299,1)]
  plot(density(g,bw="SJ"),ylim=c(0,1.2),xlim=c(0,6))
  Sys.sleep(1/30)
}
```

- Computation is often done with equally spaced bins and fast Fourier transforms.
- Methods that adjust bandwidth locally can be used.
- Some of these methods are based on nearest-neighbor fits and local polynomial fits.
- Spline based methods can be used on the log scale; the `logspline` package implements one approach.



## Density Estimation in Higher Dimensions

- Kernel density estimation can in principle be used in any number of dimensions.
- Usually a  $d$ -dimensional kernel  $K_d$  of the product form

$$K_d(u) = \prod_{i=1}^d K_1(u_i)$$

is used.

- The kernel density estimate is then

$$\hat{f}_n(x) = \frac{1}{n \det(H)} \sum_{i=1}^n K(H^{-1}(x - x_i))$$

for some matrix  $H$ .

- Suppose  $H = hA$  where  $\det(A) = 1$ . The asymptotic mean integrated square error is of the form

$$\text{AMISE} = \frac{R(K)}{nh^d} + \frac{h^4}{4} \int (\text{trace}(AA^T \nabla^2 f(x)))^2 dx$$

and therefore the optimal bandwidth and AMISE are of the form

$$h_0 = O(n^{-1/(d+4)})$$

$$\text{AMISE}_0 = O(n^{-4/(d+4)})$$

- Convergence is very slow if  $d$  is more than 2 or 3 since most of higher dimensional space will be empty—this is known as the *curse of dimensionality*.
- Density estimates in two dimensions can be visualized using perspective plots, surface plots, image plots, and contour plots.
- Higher dimensional estimates can often only be visualized by conditioning, or slicing.
- The `kde2d` function in package `MASS` provides two-dimensional kernel density estimates; an alternative is `bkde2D` in package `KernSmooth`.
- The `kde3d` function in the `misc3d` package provides three-dimensional kernel density estimates.




## Visualizing Density Estimates


Some examples are given in `geyser.R` and `kd3.R` in

<http://www.stat.uiowa.edu/~luke/classes/STAT7400-2020/examples/>

- Animation can be a useful way of understanding the effect of smoothing parameter choice.
- Bootstrap animation can help in visualizing uncertainty.
- For 2D estimates, options include
  - perspective plots
  - contour plots
  - image plots, with or without contours
- For 3D estimates contour plots are the main option
- Example: Data and contours for mixture of three trivariate normals and two bandwidths



BW = 0.2



BW = 0.5

## Kernel Smoothing and Local Regression

- A simple non-parametric regression model is

$$Y_i = m(x_i) + \varepsilon_i$$

with  $m$  a smooth mean function.

- A kernel density estimator of the conditional density  $f(y|x)$  is

$$\hat{f}_n(y|x) = \frac{\frac{1}{nh^2} \sum K\left(\frac{x-x_i}{h}\right) K\left(\frac{y-y_i}{h}\right)}{\frac{1}{nh} \sum K\left(\frac{x-x_i}{h}\right)} = \frac{1}{h} \frac{\sum K\left(\frac{x-x_i}{h}\right) K\left(\frac{y-y_i}{h}\right)}{\sum K\left(\frac{x-x_i}{h}\right)}$$

- Assuming  $K$  has mean zero, an estimate of the conditional mean is

$$\begin{aligned} \hat{m}_n(x) &= \int y \hat{f}_n(y|x) dy = \frac{\sum K\left(\frac{x-x_i}{h}\right) \int y \frac{1}{h} K\left(\frac{y-y_i}{h}\right) dy}{\sum K\left(\frac{x-x_i}{h}\right)} \\ &= \frac{\sum K\left(\frac{x-x_i}{h}\right) y_i}{\sum K\left(\frac{x-x_i}{h}\right)} = \sum w_i(x) y_i \end{aligned}$$

This is the *Nadaraya-Watson* estimator.

- This estimator can also be viewed as the result of a *locally constant* fit:  $\hat{m}_n(x)$  is the value  $\beta_0$  that minimizes

$$\sum w_i(x) (y_i - \beta_0)^2$$

- Higher degree local polynomial estimators estimate  $m(x)$  by minimizing

$$\sum w_i(x) (y_i - \beta_0 - \beta_1(x - x_i) - \cdots - \beta_p(x - x_i)^p)^2$$

- Odd values of  $p$  have advantages, and  $p = 1$ , local linear fitting, generally works well.
- Local cubic fits,  $p = 3$ , are also used.
- Problems exist near the boundary; these tend to be worse for higher degree fits.

- Bandwidth can be chosen globally or locally.
- A common local choice uses a fraction of nearest neighbors in the  $x$  direction.
- Automatic choices can use estimates of  $\sigma$  and function roughness and plug in to asymptotic approximate mean square errors.
- Cross-validation can also be used; it often undersmooths.
- Autocorrelation creates an identifiability problem.
- Software available in R includes
  - `ksmooth` for compatibility with S (but much faster).
  - `locpoly` for fitting and `dpill` for bandwidth selection in package `KernSmooth`.
  - `lowess` and `loess` for nearest neighbor based methods; also try to robustify.
  - `supsmu`, Friedman's *super smoother*, a very fast smoother.
  - package `locfit` on CRAN

All of these are also available for R; some are available as stand-alone code.

## Spline Smoothing

- Given data  $(x_1, y_1), \dots, (x_n, y_n)$  with  $x_i \in [a, b]$  one way to fit a smooth mean function is to choose  $m$  to minimize

$$S(m, \lambda) = \sum (y_i - m(x_i))^2 + \lambda \int_a^b m''(u)^2 du$$

The term  $\lambda \int_a^b m''(u)^2 du$  is a *roughness penalty*.

- Among all twice continuously differentiable functions on  $[a, b]$  this is minimized by a *natural cubic spline* with *knots* at the  $x_i$ . This minimizer is called a *smoothing spline*.
- A *cubic spline* is a function  $g$  on an interval  $[a, b]$  such that for some *knots*  $t_i$  with  $a = t_0 < t_1 < \dots < t_{n+1} = b$ 
  - on  $(t_{i-1}, t_i)$  the function  $g$  is a cubic polynomial
  - at  $t_1, \dots, t_n$  the function values, first and second derivatives are continuous.
- A cubic spline is *natural* if the second and third derivatives are zero at  $a$  and  $b$ .
- A natural cubic spline is linear on  $[a, t_1]$  and  $[t_n, b]$ .
- For a given  $\lambda$  the smoothing spline is a linear estimator.
- The set of equations to be solved is large but banded.
- The fitted values  $\hat{m}_n(x_i, \lambda)$  can be viewed as

$$\hat{m}_n(x, \lambda) = A(\lambda)y$$


where  $A(\lambda)$  is the *smoothing matrix* or *hat matrix* for the linear fit.

- The function `smooth.spline` implements smoothing splines in R.

## Example: Old Faithful Eruptions

- A nonparametric fit of waiting time to previous duration may be useful in predicting the time of the next eruption.
- The different smoothing methods considered produce the following:

```
with(geyser2, {  
  plot(pduration, waiting)  
  lines(lowess(pduration, waiting), col="red")  
  lines(supsmu(pduration, waiting), col="blue")  
  lines(ksmooth(pduration, waiting), col="green")  
  lines(smooth.spline(pduration, waiting), col="orange")  
})
```



- An animated version of the smoothing spline (available on line) shows the effect of varying the smoothing parameter.



## Degrees of Freedom of a Linear Smoother

- For a linear regression fit with hat matrix

$$H = X(X^T X)^{-1} X^T$$

and full rank regressor matrix  $X$

$\text{tr}(H)$  = number of fitted parameters = degrees of freedom of fit

- By analogy define the degrees of freedom of a linear smoother as

$$\text{df}_{\text{fit}} = \text{tr}(A(\lambda))$$

For the geyser data, the degrees of freedom of a smoothing spline fit with the default bandwidth selection rule are

```
> sum(with(geyser2, smooth.spline(pduraction, waiting))$lev)
[1] 4.169843
> with(geyser2, smooth.spline(pduraction, waiting))$df
[1] 4.169843
```

- For residual degrees of freedom the definition usually used is

$$\text{df}_{\text{res}} = n - 2\text{tr}(A(\lambda)) + \text{tr}(A(\lambda)A(\lambda)^T)$$

- Assuming constant error variance, a possible estimate is

$$\hat{\sigma}_{\varepsilon}^2 = \frac{\sum (y_i - \hat{m}_n(x_i, \lambda))^2}{\text{df}_{\text{res}}(\lambda)} = \frac{\text{RSS}(\lambda)}{\text{df}_{\text{res}}(\lambda)}$$

- The simpler estimator

$$\hat{\sigma}_{\varepsilon}^2 = \frac{\text{RSS}(\lambda)}{\text{tr}(I - A(\lambda))} = \frac{\text{RSS}(\lambda)}{n - \text{df}_{\text{fit}}}$$

is also used.

- To reduce bias it may make sense to use a rougher smooth for variance estimation than for mean function estimation.

## Choosing Smoothing Parameters for Linear Smoothers

- Many smoothing methods are linear for a given value of a smoothing parameter  $\lambda$ .
- Choice of the smoothing parameter  $\lambda$  can be based on leave-one-out cross-validation, i.e. minimizing the *cross-validation score*

$$\text{CV}(\lambda) = \frac{1}{n} \sum (y_i - \hat{m}_n^{(-i)}(x_i, \lambda))^2$$

- If the smoother satisfies (at least approximately)

$$\hat{m}_n^{(-i)}(x_i, \lambda) = \frac{\sum_{j \neq i} A(\lambda)_{ij} y_j}{\sum_{j \neq i} A(\lambda)_{ij}}$$

and

$$\sum_{j=1}^n A(\lambda)_{ij} = 1 \quad \text{for all } i$$

then the cross-validation score can be computed as

$$\text{CV}(\lambda) = \frac{1}{n} \sum \left( \frac{y_i - \hat{m}_n(x_i, \lambda)}{1 - A_{ii}(\lambda)} \right)^2$$

- The *generalized cross-validation criterion*, or GCV, uses average leverage values:

$$\text{GCV}(\lambda) = \frac{1}{n} \sum \left( \frac{y_i - \hat{m}_n(x_i, \lambda)}{1 - n^{-1} \text{trace}(A(\lambda))} \right)^2$$


- The original motivation for GCV was computational; with better algorithms this is no longer an issue.

- An alternative motivation for GCV:
  - For an orthogonal transformation  $Q$  one can consider fitting  $y_Q = QY$  with  $A_Q(\lambda) = QA(\lambda)Q^T$ .
  - Coefficient estimates and  $SS_{\text{res}}$  are the same for all  $Q$ , but the CV score is not.
  - One can choose an orthogonal transformation such that the diagonal elements of  $A_Q(\lambda)$  are constant.
  - For any such  $Q$  we have  $A_Q(\lambda)_{ii} = n^{-1}\text{trace}(A_Q(\lambda)) = n^{-1}\text{trace}(A(\lambda))$
- Despite the name, GCV does not generalize CV.
- Both CV and GCV have a tendency to undersmooth.

- For the geyser data the code

```
with(geyser2, {
  lambda <- seq(0.5, 2, len=30)
  f <- function(s, cv = FALSE)
    smooth.spline(pduraction, waiting, spar=s, cv=cv)$cv
  gcv <- sapply(lambda, f)
  cv <- sapply(lambda, f, TRUE)
  plot(lambda, gcv, type="l")
  lines(lambda, cv, col="blue")
})
```

extracts and plots GCV and CV values:



- Both criteria select a value of  $\lambda$  close to 1.

- Other smoothing parameter selection criteria include

- Mallows  $C_p$ ,

$$C_p = \text{RSS}(\lambda) + 2\hat{\sigma}_\varepsilon^2 \text{df}_{\text{fit}}(\lambda)$$

- Akaike's information criterion (AIC)

$$\text{AIC}(\lambda) = \log\{\text{RSS}(\lambda)\} + 2\text{df}_{\text{fit}}(\lambda)/n$$

- Corrected AIC of Hurvich, Simonoff, and Tsai (1998)

$$\text{AIC}_C(\lambda) = \log\{\text{RSS}(\lambda)\} + \frac{2(\text{df}_{\text{fit}}(\lambda) + 1)}{n - \text{df}_{\text{fit}}(\lambda) - 2}$$

## Spline Representations

- Splines can be written in terms of many different bases,
  - B-splines
  - truncated power basis
  - radial or thin plate basis

Some are more useful numerically, others have interpretational advantages.

- One useful basis for a cubic spline with knots  $\{\kappa_1, \dots, \kappa_K\}$  is the *radial basis* or *thin plate basis*

$$1, x, |x - \kappa_1|^3, \dots, |x - \kappa_K|^3$$

- More generally, a basis for splines of order  $2m - 1$  is

$$1, x, \dots, x^{m-1}, |x - \kappa_1|^{2m-1}, \dots, |x - \kappa_K|^{2m-1}$$

for  $m = 1, 2, 3, \dots$

- $m = 2$  produces cubic splines
  - $m = 1$  produces linear splines
- In terms of this basis a spline is a function of the form

$$f(x) = \sum_{j=0}^{m-1} \beta_j x^j + \sum_{k=1}^K \delta_k |x - \kappa_k|^{2m-1}$$

- References:
  - P. J. Green and B. W. Silverman (1994). *Nonparametric Regression and Generalized Linear Models*
  - D. Ruppert, M. P. Wand, and R. J. Carroll (2003). *Semiparametric Regression*. `SemiPar` is an R package implementing the methods of this book.
  - G. Wahba (1990). *Spline Models for Observational Data*.
  - S. Wood (2017). *Generalized Additive Models: An Introduction with R, 2nd Ed.*. This is related to the `mgcv` package.

- A generic form for the fitted values is

$$\hat{y} = X_0\beta + X_1\delta.$$

- *Regression splines* refers to models with a small number of knots  $K$  fit by ordinary least squares, i.e. by choosing  $\beta, \delta$  to minimize

$$\|y - X_0\beta - X_1\delta\|^2$$

- *Penalized spline smoothing* fits models with a larger number of knots subject to a quadratic constraint

$$\delta^T D \delta \leq C$$

for a positive definite  $D$  and some  $C$ .

- Equivalently, by a Lagrange multiplier argument, the solution minimizes the penalized least squares criterion

$$\|y - X_0\beta - X_1\delta\|^2 + \lambda \delta^T D \delta$$

for some  $\lambda > 0$ .

- A common form of  $D$  is

$$D = \left[ |\kappa_i - \kappa_j|^{2m-1} \right]_{1 \leq i, j \leq K}$$

- A variant uses

$$D = \Omega^{1/2} (\Omega^{1/2})^T$$

with

$$\Omega = \left[ |\kappa_i - \kappa_j|^{2m-1} \right]_{1 \leq i, j \leq K}$$

where the *principal square root*  $M^{1/2}$  of a matrix  $M$  with SVD

$$M = U \text{diag}(d) V^T$$

is defined as

$$M^{1/2} = U \text{diag}(\sqrt{d}) V^T$$

This form ensures that  $D$  is at least positive semi-definite.

- Smoothing splines are penalized splines of degree  $2m - 1 = 3$  with knots  $\kappa_i = x_i$  and

$$D = [|\kappa_i - \kappa_j|^3]_{1 \leq i, j \leq n}$$

and the added natural boundary constraint

$$X_0^T \delta = 0$$

- For a natural cubic spline

$$\int g''(t)^2 dt = \delta^T D \delta$$

The quadratic form  $\delta^T D \delta$  is strictly positive definite on the subspace defined by  $X_0^T \delta = 0$ .

- Penalized splines can often approximate smoothing splines well using far fewer knots.
- The detailed placement of knots and their number is usually not critical as long as there are enough.
- Simple default rules that often work well (Ruppert, Wand, and Carroll 2003):

– knot locations:

$$\kappa_k = \left( \frac{k+1}{K+2} \right) \text{th sample quantile of unique } x_i$$

– number of knots:

$$K = \min \left( \frac{1}{4} \times \text{number of unique } x_i, 35 \right)$$

The `SemiPar` package actually seems to use the default

$$K = \max \left( \frac{1}{4} \times \text{number of unique } x_i, 20 \right)$$

- More sophisticated methods for choosing number and location of knots are possible but not emphasized in the penalized spline literature at this point.



## A Useful Computational Device

To minimize

$$\|Y - X_0\beta - X_1\delta\|^2 + \lambda \delta^T D \delta$$

for a given  $\lambda$ , suppose  $B$  satisfies

$$\lambda D = B^T B$$

and

$$Y^* = \begin{bmatrix} Y \\ 0 \end{bmatrix} \quad X^* = \begin{bmatrix} X_0 & X_1 \\ 0 & B \end{bmatrix} \quad \beta^* = \begin{bmatrix} \beta \\ \delta \end{bmatrix}$$

Then

$$\|Y^* - X^* \beta^*\|^2 = \|Y - X_0\beta - X_1\delta\|^2 + \lambda \delta^T D \delta$$

So  $\hat{\beta}$  and  $\hat{\delta}$  can be computed by finding the OLS coefficients for the regression of  $Y^*$  on  $X^*$ .

## Penalized Splines and Mixed Models

- For strictly positive definite  $D$  and a given  $\lambda$  minimizing the objective function

$$\|y - X_0\beta - X_1\delta\|^2 + \lambda \delta^T D \delta$$

is equivalent to maximizing the log likelihood for the mixed model

$$Y = X_0\beta + X_1\delta + \varepsilon$$

with fixed effects parameters  $\beta$  and

$$\varepsilon \sim N(0, \sigma_\varepsilon^2 I)$$

$$\delta \sim N(0, \sigma_\delta^2 D^{-1})$$

$$\lambda = \sigma_\varepsilon^2 / \sigma_\delta^2$$

with  $\lambda$  known.

- Some consequences:
  - The penalized spline fit at  $x$  is the BLUP for the mixed model with known mixed effects covariance structure.
  - Linear mixed model software can be used to fit penalized spline models (the R package `SemiPar` does this).
  - The smoothing parameter  $\lambda$  can be estimated using ML or REML estimates of  $\sigma_\varepsilon^2$  and  $\sigma_\delta^2$  from the linear mixed model.
  - Interval estimation/testing formulations from mixed models can be used.
- Additional consequences:
  - The criterion has a Bayesian interpretation.
  - Extension to models containing smoothing and mixed effects are immediate.
  - Extension to generalized linear models can use GLMM methodology.

## Example: Old Faithful Eruptions

- Using the function `spm` from `SemiPar` a penalized spline model can be fit with


```
> library(SemiPar)
> attach(geyser2) # needed because of flaws in spm implementation
> summary(spm(waiting ~ f(pduration)))
Summary for non-linear components:
```

```
              df spar knots
f(pduration) 4.573  2.9   28
```

Note this includes 1 df for the intercept.

- The `plot` method for the `spm` result produces a plot with pointwise error bars:

```
> plot(spm(waiting ~ f(pduration)), ylim = range(waiting))
> points(pduration, waiting)
```



**A fit using mgcv:**

```
> library(mgcv)
> gam.fit <- gam(waiting ~ s(pduration), data = geyser2)
> summary(gam.fit)
```

```
Family: gaussian
Link function: identity
```

```
Formula:
waiting ~ s(pduration)
```

```
Parametric coefficients:
```

	Estimate	Std. Error	t value	Pr(> t )
(Intercept)	72.2886	0.3594	201.1	<2e-16 ***

```
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
Approximate significance of smooth terms:
```

	edf	Ref.df	F	p-value
s(pduration)	3.149	3.987	299.8	<2e-16 ***

```
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
R-sq.(adj) = 0.801    Deviance explained = 80.3%
GCV = 39.046   Scale est. = 38.503      n = 298
```

**A plot of the smooth component with the mean-adjusted waiting times is produced by**

```
> plot(gam.fit)
> with(geyser2, points(pduration, waiting - mean(waiting)))
```

## Smoothing with Multiple Predictors

- Many methods have natural generalizations
- All suffer from the curse of dimensionality.
- Generalizations to two or three variables can work reasonably.
- Local polynomial fits can be generalized to  $p$  predictors.
- `loess` is designed to handle multiple predictors, in principle at least.
- Spline methods can be generalized in two ways:
  - *tensor product splines* use all possible products of single variable spline bases.
  - *thin plate splines* generalize the radial basis representation.
- A thin plate spline of order  $m$  in  $d$  dimensions is of the form

$$f(x) = \sum_{i=1}^M \beta_i \phi_i(x) + \sum_{k=1}^K \delta_k r(x - \kappa_k)$$

with

$$r(u) = \begin{cases} \|u\|^{2m-d} & \text{for } d \text{ odd} \\ \|u\|^{2m-d} \log \|u\| & \text{for } d \text{ even} \end{cases}$$

and where the  $\phi_i$  are a basis for the space of polynomials of total degree  $\leq m-1$  in  $d$  variables. The dimension of this space is

$$M = \binom{d+m-1}{d}$$

If  $d=2, m=2$  then  $M=3$  and a basis is

$$\phi_1(x) = 1, \phi_2(x) = x_1, \phi_3(x) = x_2$$

## Penalized Thin Plate Splines

- Penalized thin plate splines usually use a penalty with

$$D = \Omega^{1/2}(\Omega^{1/2})^T$$

where

$$\Omega = [r(\kappa_i - \kappa_j)]_{1 \leq i, j \leq K}$$

This corresponds at least approximately to using a squared derivative penalty.

- Simple knot selection rules are harder for  $d > 1$ .
- Some approaches:
  - space-filling designs (Nychka and Saltzman, 1998)
  - clustering algorithms, such as `clara`

## Multivariate Smoothing Splines

- The bivariate smoothing spline objective of minimizing

$$\sum (y_i - g(x_i))^2 + \lambda J(g)$$

with

$$J(g) = \int \int \left( \frac{\partial^2 g}{\partial x_1^2} \right)^2 + 2 \left( \frac{\partial^2 g}{\partial x_1 \partial x_2} \right)^2 + \left( \frac{\partial^2 g}{\partial x_2^2} \right)^2 dx_1 dx_2$$

is minimized by a thin plate spline with knots at the  $x_i$  and a constraint on the  $\delta_k$  analogous to the natural spline constraint.

- Scaling of variables needs to be addressed
- Thin-plate spline smoothing is closely related to *kriging*.
- The general smoothing spline uses

$$D = X_1 = [r(\kappa_i - \kappa_i)]$$

with the constraint  $X_0^T \delta = 0$ .

- Challenge: the linear system to be solved for each  $\lambda$  value to fit a smoothing spline is large and not sparse.

## Thin Plate Regression Splines

- Wood (2017) advocates an approach called *thin plate regression splines* that is implemented in the `mgcv` package.
- The approach uses the spectral decomposition of  $X_1$

$$X_1 = UEU^T$$

with  $E$  the diagonal matrix of eigen values, and the columns of  $U$  the corresponding eigen vectors.

- The eigen values are ordered so that  $|E_{ii}| \geq |E_{jj}|$  for  $i \leq j$ .
- The approach replaces  $X_1$  with a lower rank approximation

$$X_{1,k} = U_k E_k U_k^T$$

using the  $k$  largest eigen values in magnitude.

- The implementation uses an iterative algorithm (Lanczos iteration) for computing the largest  $k$  eigenvalues/singular values and vectors.
- The  $k$  leading eigenvectors form the basis for the fit.
- The matrix  $X_1$  does not need to be formed explicitly; it is enough to be able to compute  $X_1 v$  for any  $v$ .
- $k$  could be increased until the change in estimates is small or a specified limit is reached.
- As long as  $k$  is large enough results are not very sensitive to the particular value of  $k$ .
- `mgcv` by default uses  $k = 10 \times 3^{d-1}$  for a  $d$ -dimensional smooth.



- This approach seems to be very effective in practice and avoids the need to specify a set of knots.
- The main drawback is that the choice of  $k$  and its impact on the basis used are less interpretable.
- With this approach the computational cost is reduced from  $O(n^3)$  to  $O(n^2k)$ .
- For large  $n$  Wood (2017) recommends using a random sample of  $n_r$  rows to reduce the computation cost to  $O(n_r^2k)$ . (From the help files the approach in `mgcv` looks more like  $O(n \times n_r \times k)$  to me).

## Example: Scallop Catches



- Data records location and size of scallop catches off Long Island.
- A bivariate penalized spline fit is computed by

```
> data(scallop)
> attach(scallop)
> log.catch <- log(tot.catch + 1)
> fit <- spm(log.catch ~ f(longitude, latitude))
> summary(fit)
```

Summary for non-linear components:

	df	spar	knots
f(longitude, latitude)	25.12	0.2904	37

- Default knot locations are determined using `clara`
- Knot locations and fit:



A fit using `mgcv` would use

```
> scallop.gam <- gam(log.catch ~ s(longitude, latitude), data = scallop)
> summary(scallop.gam)
```

Family: gaussian  
Link function: identity

Formula:  
log.catch ~ s(longitude, latitude)

Parametric coefficients:

	Estimate	Std. Error	t value	Pr(> t )
(Intercept)	3.4826	0.1096	31.77	<2e-16 ***

---  
Signif. codes: 0 '\*\*\*' 0.001 '\*\*' 0.01 '\*' 0.05 '.' 0.1 ' ' 1

Approximate significance of smooth terms:

	edf	Ref.df	F	p-value
s(longitude,latitude)	26.23	28.53	8.823	<2e-16 ***

---  
Signif. codes: 0 '\*\*\*' 0.001 '\*\*' 0.01 '\*' 0.05 '.' 0.1 ' ' 1

R-sq.(adj) = 0.623    Deviance explained = 69%  
GCV = 2.1793    Scale est. = 1.7784    n = 148  
> plot(scallop.gam)

## Computational Issues

- Algorithms that select the smoothing parameter typically need to compute smooths for many parameter values.
- Smoothing splines require solving an  $n \times n$  system.
  - For a single variable the fitting system can be made tri-diagonal.
  - For thin plate splines of two or more variables the equations are not sparse.
- Penalized splines reduce the computational burden by choosing fewer knots, but then need to select knot locations.
- Thin plate regression splines (implemented in the `mgcv` package) use a rank  $k$  approximation for a user-specified  $k$ .
- As long as the number of knots or the number of terms  $k$  is large enough results are not very sensitive to the particular value of  $k$ .
- Examples are available in

<http://www.stat.uiowa.edu/~luke/classes/STAT7400-2020/examples/smoothex.Rmd>