

A Duality Proof of Tchakaloff's Theorem^{*†}

Ral E. Curto

The University of Iowa
E-mail: curto@math.uiowa.edu

and

Lawrence A. Fialkow[‡]

State University of New York, New Paltz
E-mail: fialkowl@newpaltz.edu

Tchakaloff's Theorem establishes the existence of a quadrature rule of prescribed degree relative to a positive, compactly supported measure that is absolutely continuous with respect to Lebesgue measure on \mathbb{R}^d . Subsequent extensions were obtained by Mysovskikh and by Putinar. We provide new proofs and partial extensions of these results, based on duality techniques utilized by Stochel. We also obtain new uniqueness criteria in the Truncated Complex Moment Problem.

Key Words: quadratures, truncated moment problems, Tchakaloff's theorem, duality

1. INTRODUCTION

Tchakaloff's Theorem [25, Théorème II] establishes the existence of a quadrature rule of prescribed degree relative to a positive, compactly supported measure that is absolutely continuous with respect to Lebesgue measure on \mathbb{R}^d . An extension to the case when the support is unbounded was subsequently obtained by Mysovskikh [15], and in [17], M. Putinar generalized these results to arbitrary positive Borel measures. In the present note we provide new proofs, and partial extensions, of Putinar's results,

^{*} 1991 *Mathematics Subject Classification*. Primary 47A57, 46A20, 65D32, 44A60; Secondary 47N40, 46B22

[†] Research partially supported by NSF grants.

[‡] Also partially supported by the State University of New York at New Paltz Research and Creative Projects Award Program.

based on duality techniques utilized by J. Stochel [22] in a recent study of multivariable moment problems. We also obtain some new uniqueness criteria in the Truncated Complex Moment Problem (cf. [5], [6], [7]).

For $t \equiv (t_1, \dots, t_d) \in \mathbb{R}^d$, and for a multi-index $i \equiv (i_1, \dots, i_d) \in \mathbb{Z}_+^d$, let $t^i = t_1^{i_1} \cdots t_d^{i_d}$ and $|i| = i_1 + \cdots + i_d$. Let $\mathbb{R}_{m,d}[t] \equiv \mathbb{R}_m[t_1, \dots, t_d]$ denote the space of real polynomials of total degree at most m in t_1, \dots, t_d , and let $N_{m,d} := \dim \mathbb{R}_{m,d}[t]$. As a notational convenience, to indicate that each function in $\mathbb{R}_{m,d}[t]$ is (absolutely) integrable with respect to a positive Borel measure μ on \mathbb{R}^d , we write $\mathbb{R}_{m,d}[t] \subseteq L^1(\mu)$; note that in this case, the canonical map from $\mathbb{R}_{m,d}[t]$ into $L^1(\mu)$ need not be one-to-one. For a measure μ with closed support $K \subseteq \mathbb{R}^d$ and satisfying $\mathbb{R}_{m,d}[t] \subseteq L^1(\mu)$, a quadrature rule of precision (or degree) m and size $N (< \infty)$ consists of nodes x_1, \dots, x_N in K and positive weights ρ_1, \dots, ρ_N such that

$$\int p(t) d\mu(t) = \sum_{k=1}^N \rho_k p(x_k) \quad (p \in \mathbb{R}_{m,d}[t]). \quad (1.1)$$

Putinar's first generalization of Tchakaloff's Theorem concerns the case of compact support.

THEOREM 1.1. [17, Theorem 1] *Let μ be a positive, finite Borel measure with compact support in \mathbb{R}^d . Given $m > 0$, there exists a quadrature rule for μ of degree m with size $\leq N_{m,d}$.*

REMARK 1.2. *For a positive Borel measure μ on \mathbb{R}^d having convergent moments up to at least degree n , let $N_{n,d;\mu} := \dim \{p|_{\text{supp } \mu} : p \in \mathbb{R}_{n,d}[t]\}$. We show in Section 3 that the estimate $N_{m,d}$ in Theorem 1.1 can be refined to $N_{m,d;\mu}$ (cf. Theorem 3.5).*

For the case of unbounded support, Putinar's quadrature result assumes a somewhat different form.

THEOREM 1.3. [17, Theorem 2] *Let μ be a positive Borel measure supported in \mathbb{R}^d , and suppose that μ has convergent moments up to at least degree $2m$, i.e., $\mathbb{R}_{2m,d}[t] \subseteq L^1(\mu)$. Then there exists a quadrature rule for μ of degree $2m - 1$ with size $\leq N_{2m,d}$.*

In [17, Corollary 1], Putinar also obtained an analogue of Theorem 1.3 valid in any degree (even or odd), provided $\text{supp } \mu$ is contained in a proper closed convex cone of \mathbb{R}^d . In our main result, which follows, we are able to remove the latter constraint on $\text{supp } \mu$, so as to treat the "even" and "odd" cases together, with an improved estimate for the size of a quadrature rule.

THEOREM 1.4. (*Generalized Tchakaloff Theorem, real case*) *Let μ be a positive Borel measure on \mathbb{R}^d having convergent moments up to at least degree n . Then there exists a quadrature rule for μ of degree $n - 1$ with size $\leq 1 + N_{n-1,d;\mu}$.*

REMARK 1.5. *For a quadrature rule of degree $n - 1$, the size estimate $1 + N_{n-1,d;\mu}$ of Theorem 1.4 compares favorably with the estimate $N_{n,d}$ of Theorem 1.3 and [17, Corollary 1]. For the case of compact support, Tchakaloff's size estimate $N_{n-1,d}$ (for a rule of degree $n - 1$) is known to be sharp [25, pp. 131-133]. It is thus plausible that $1 + N_{n-1,d;\mu}$ is sharp in the non-compact case, though we do not have an example confirming this. Of course, for certain sets and measures there exist Gaussian-type quadrature rules for which the size is much smaller than that guaranteed by the above estimates [14], [18], [20], [21], [24], [27]. In particular, for $d = 1$, any positive Borel measure on \mathbb{R} , $[a, b]$, or $[0, +\infty)$, having moments up to degree n , admits a Gaussian-type quadrature rule of degree n with size $\leq \lfloor \frac{n}{2} \rfloor + 1$ [11].*

As we next describe, Theorems 1.1, 1.3 and 1.4 are closely related to the following Truncated Multivariable Moment Problem (TMMP) [5, Chapter 7]. For a real multisequence $\beta \equiv \beta^{(m)} = \{\beta_i\}_{i \in \mathbb{Z}_+^d, |i| \leq m}$, TMMP seeks to characterize the existence of a positive Borel measure μ supported in \mathbb{R}^d such that

$$\beta_i = \int t^i d\mu(t) \quad (|i| \leq m); \quad (1.2)$$

a measure μ as in (1.2) is said to be a *representing measure* for β . The *K-moment problem* further requires $\text{supp } \mu \subseteq K$, where K is a prescribed closed subset of \mathbb{R}^d . The following basic question remains open.

QUESTION 1.6. *If $\beta \equiv \beta^{(m)}$ has a representing measure, does β admit a finitely atomic representing measure (i.e., a measure of the form $\sum_{k=1}^N \rho_k \delta_{x_k}$, where $1 \leq N < \infty$, each $\rho_k > 0$, and δ_{x_k} is the point mass at $x_k \in \mathbb{R}^d$)? More generally, if β has a representing measure supported in K , does β have a finitely atomic representing measure supported in K ?*

Let μ be a representing measure for $\beta \equiv \beta^{(m)}$ and let $K := \text{supp } \mu$. The existence of a finitely atomic representing measure in the *K-moment problem* for β is equivalent to the existence of a quadrature rule for μ of degree m . For $d = 1$ and $K = \mathbb{R}$, $[a, b]$, or $[0, \infty)$, [11] implies that

Question 1.6 has an affirmative answer relative to representing measures in the K -moment problem. For $d \geq 1$, Theorem 1.1 shows that if β has a representing measure supported in a compact set K , then β admits a finitely atomic representing measure supported in K . Theorem 1.4 implies that if $\beta \equiv \beta^{(m)}$ has a representing measure μ satisfying $\mathbb{R}_{m+1,d}[t] \subseteq L^1(\mu)$ and $\text{supp } \mu \subseteq K$ (closed), then β admits a finitely atomic representing measure supported in K .

The proof of Theorem 1.3 in [17] rests on convexity arguments similar to those in [25], and also on some rather subtle convergence arguments. Our proof of Theorem 1.4 is based on a different approach. Given a multisequence $\beta \equiv \beta^{(m)}$ and a prescribed closed set $K \subseteq \mathbb{R}^d$, let $M_K(\beta)$ denote the set of representing measures for β supported in K . In Proposition 2.3, we show that any extreme point of $M_K(\beta)$ is finitely atomic. This result is based on a technique of J.P. Gabardo [13], who attributes the underlying idea to Naimark (cf. [1]). Proposition 2.3 yields some insight into Question 1.6; in particular, if there is a unique representing measure for β , it is finitely atomic. Let μ be a measure satisfying the hypothesis of Theorem 1.4, i.e., $\mathbb{R}_{n,d}[t] \subseteq L^1(\mu)$. In Section 3 we employ duality results of Stochel [22] to show that a certain convex set of representing measures for $\beta^{(n-1)}[\mu]$ admits an extreme point, which then acts as a quadrature rule for μ of degree $n - 1$.

In Section 4 we present some uniqueness results in the following Truncated Complex Moment Problem (TCMP) [5]. Let $\gamma \equiv \gamma^{(2n)} : \gamma_{00}, \gamma_{01}, \gamma_{10}, \gamma_{02}, \gamma_{11}, \gamma_{20}, \dots, \gamma_{0,2n}, \dots, \gamma_{2n,0}$ denote a sequence of complex numbers. TCMP entails characterizing the existence of a positive Borel measure μ on \mathbb{C} such that

$$\gamma_{ij} = \int \bar{z}^i z^j d\mu(z) \quad (0 \leq i + j \leq 2n).$$

We encode the data γ in a moment matrix $M(n)(\gamma)$ [5, Chapter 2], whose successive columns are labeled $1, Z, \bar{Z}, \dots, Z^n, \dots, \bar{Z}^n$. If γ has a representing measure, then $M(n)(\gamma)$ is positive semidefinite and *recursively generated* (cf. [5, (3.2) and Remark 3.15(ii)]); moreover, if μ is a representing measure for γ , then $\text{card supp } \mu \geq \text{rank } M(n)$ [5, Corollary 3.7]. In [5, Chapter 5], for the case of *flat data*, where $M(n) \geq 0$ and $\text{rank } M(n) = \text{rank } M(n-1)$, we established the existence of (and explicitly constructed) a unique finitely atomic representing measure. In Proposition 4.1 we show that this measure (with $\text{rank } M(n)$ atoms) is actually the unique representing measure for γ .

We say that $M(n)(\gamma)$ admits an *analytic relation* if there exist $k \leq n$ and scalars $a_{ij} \in \mathbb{C}$ ($0 \leq i + j < k$) such that in the column space of $M(n)$

there is a dependence relation

$$Z^k = \sum_{0 \leq i+j < k} a_{ij} \bar{Z}^i Z^j. \quad (1.3)$$

In [6, Theorem 3.1] we proved that if $M(n)$ is positive and recursively generated, and if $M(n)$ admits an analytic relation with $k \leq \lfloor \frac{n}{2} \rfloor + 1$, then γ admits a unique finitely atomic representing measure (with rank $M(n)$ atoms); by contrast, for $k > \lfloor \frac{n}{2} \rfloor + 1$, there need not be any representing measure [7, Example 2.14], or there may be a finitely atomic representing measure, but none with as few as rank $M(n)$ atoms [12, Theorem 3.1]. In Proposition 4.2 we prove that if γ has a representing measure and $M(n)(\gamma)$ admits an analytic relation (as in (1.3)) for some $k \leq n$, then γ has a unique representing measure, which is finitely atomic, with at most k^2 atoms. This result depends on the fact that a polynomial of the form $z^k - q(z, \bar{z})$ ($\deg q < k$) has at most k^2 roots (Proposition 4.4).

2. EXTREME POINTS AND FINITELY ATOMIC REPRESENTING MEASURES

For $z \equiv (z_1, \dots, z_d) \in \mathbb{C}^d$ and $j \equiv (j_1, \dots, j_d) \in \mathbb{Z}_+^d$, let $z^j = z_1^{j_1} \cdots z_d^{j_d}$ and let $|j| = j_1 + \cdots + j_d$. Let $\mathbb{C}_{m,d}[z, \bar{z}]$ denote the complex polynomials $p(z, \bar{z})$ of total degree at most m . For a closed set $K \subseteq \mathbb{C}^d$, $\mathbb{C}_{m,d}[z, \bar{z}]|_K$ denotes the vector space of restrictions to K of polynomials in $\mathbb{C}_{m,d}[z, \bar{z}]$. Given a complex sequence $\gamma \equiv \gamma^{(m)} = \{\gamma_{ij}\}_{i,j \in \mathbb{Z}_+^d, |i|+|j| \leq m}$, and a closed set $K \subseteq \mathbb{C}^d$, the Multivariable Truncated Complex K -Moment Problem entails characterizing the existence of a positive Borel measure μ , supported in K , such that

$$\gamma_{ij} = \int \bar{z}^i z^j d\mu, \quad |i| + |j| \leq m$$

(where $\bar{z}^i = \bar{z}_1^{i_1} \cdots \bar{z}_d^{i_d}$). The Multivariable *Full* Complex K -Moment Problem in \mathbb{C}^d concerns the analogous problem, for a sequence $\gamma^{(\infty)} \equiv \{\gamma_{i,j}\}_{i,j \in \mathbb{Z}_+^d}$, which prescribes moments of all orders. The Full and Truncated Complex K -Moment Problems on \mathbb{C}^d are equivalent, respectively, to corresponding moment problems on \mathbb{R}^{2d} (cf. [5, Chapters 6 and 7], [8, Section 5], [9] and [23, Appendix]). The following result of J. Stochel provides the connection between the Full and Truncated Multivariable Complex K -Moment Problems.

THEOREM 2.1. [22, Theorem 4] *$\gamma^{(\infty)}$ has a representing measure supported in a closed set $K \subseteq \mathbb{C}^d$ if and only if, for each $m > 0$, $\gamma^{(m)}$ admits a representing measure supported in K .*

Assume that $\gamma \equiv \gamma^{(m)}$ admits a representing measure supported in a closed set $K \subseteq \mathbb{C}^d$. Consider the convex set

$$M_K(\gamma) = \{\nu : \nu \text{ is a representing measure for } \gamma \text{ and } \text{supp } \nu \subseteq K\}.$$

It is not known whether $M_K(\gamma)$ always has an extreme point. The proof of the following result is motivated by an argument in [13, Proposition 2.5 and Corollary 2.6]. For $\nu \in M_K(\gamma)$, let $\mathbb{C}_{m,d}[z, \bar{z}](\nu)$ denote the image of $\mathbb{C}_{m,d}[z, \bar{z}]$ in $L^1(\nu)$ under the canonical projection.

PROPOSITION 2.2. *If $M_K(\gamma)$ has an extreme point ν , then ν is finitely atomic, with $\text{card supp } \nu \leq \dim \mathbb{C}_{m,d}[z, \bar{z}](\nu) (\leq \dim \mathbb{C}_{m,d}[z, \bar{z}]|_K)$.*

Proof. Since ν is a representing measure for $\gamma^{(m)}$, we may consider $\mathcal{L} := \mathbb{C}_{m,d}[z, \bar{z}](\nu) \subseteq L^1(\nu)$, and we claim that \mathcal{L} is dense in $L^1(\nu)$.

Since ν is finite, it follows that $L^1(\nu)^* = L^\infty(\nu)$ [3, Theorem III.5.6]; thus, if \mathcal{L} is not dense, there exists $f \in L^\infty(\nu)$, $f \neq 0$, such that

$$\int pf \, d\nu = 0 \quad (p \in \mathcal{L}). \quad (2.2.1)$$

Since $\nu \geq 0$ and \mathcal{L} is self-adjoint, we may replace f by $\frac{1}{2\|f+\bar{f}\|_\infty} (f+\bar{f})$, and we may thus assume that f is real, with $\|f\|_\infty \leq \frac{1}{2}$. Thus $\nu_1 := (1+f)\nu$ and $\nu_2 := (1-f)\nu$ are positive, and (2.2.1) implies that they belong to $M_K(\gamma)$. Since $\nu = \frac{1}{2}\nu_1 + \frac{1}{2}\nu_2$, we have a contradiction to the hypothesis that ν is an extreme point of $M_K(\gamma)$.

Now \mathcal{L} is dense in $L^1(\nu)$, and since \mathcal{L} is finite dimensional, we have $\mathcal{L} = L^1(\nu)$, whence $r := \dim \mathcal{L} = \dim L^1(\nu) = \dim L^1(\nu)^* = \dim L^\infty(\nu)$. Suppose $\text{supp } \nu$ contains distinct points z_1, \dots, z_{r+1} , and let $\{D_i\}_{i=1}^{r+1}$ denote mutually disjoint closed disks of positive radii such that $z_i \in D_i$ ($1 \leq i \leq r+1$). Then $\{\chi_{D_i}\}_{i=1}^{r+1}$ is linearly independent in $L^\infty(\nu)$; this contradiction implies

$$\begin{aligned} \text{card supp } \nu &\leq r = \dim \mathbb{C}_{m,d}[z, \bar{z}](\nu) \\ &\leq \dim \mathbb{C}_{m,d}[z, \bar{z}]|_{\text{supp } \nu} \leq \dim \mathbb{C}_{m,d}[z, \bar{z}]|_K. \end{aligned}$$

■

The analogue of Proposition 2.2 for a real multisequence $\beta^{(m)}$ and a closed subset $K \subseteq \mathbb{R}^d$ can be proved by a straightforward modification of the preceding argument; we omit the details.

PROPOSITION 2.3. *If $M_K(\beta^{(m)})$ has an extreme point ν , then ν is finitely atomic, with $\text{card supp } \nu \leq \dim \mathbb{R}_{m,d}[t](\nu) (\leq \dim \mathbb{R}_{m,d}[t] |_K)$.*

3. A DUALITY PROOF OF TCHAKALOFF'S THEOREM

The main result of this section is the following complex version of the generalized Tchakaloff theorem; we treat the complex case first mostly as a convenience, since the tools we require from [22] are formulated in terms of the complex moment problem.

For a positive Borel measure μ on \mathbb{C}^d having convergent moments up to at least degree n , recall that $\mathbb{C}_{n,d}[z, \bar{z}](\mu)$ denotes the image of $\mathbb{C}_{n,d}[z, \bar{z}]$ in $L^1(\mu)$ under the canonical projection, and let $\mathcal{N}_{n,d}(\mu) := \dim \mathbb{C}_{n,d}[z, \bar{z}](\mu)$ and $N_{n,d;\mu} := \dim \mathbb{C}_{n,d}[z, \bar{z}] |_{\text{supp } \mu}$.

THEOREM 3.1. *(Generalized Tchakaloff Theorem, complex case). Let μ be a positive Borel measure on \mathbb{C}^d having convergent moments up to at least degree n , and let $K := \text{supp } \mu$. Then, for some $N \leq 1 + N_{n-1,d;\mu}$, there exist nodes $z_1, \dots, z_N \in K$, and positive weights ρ_1, \dots, ρ_N , such that*

$$\int_K p(z, \bar{z}) d\mu(z) = \sum_{k=1}^N \rho_k p(z_k, \bar{z}_k) \quad (p \in \mathbb{C}_{n-1,d}[z, \bar{z}]).$$

To prove Theorem 3.1, we first introduce some preliminary results and notation concerning duality. Let X be a locally compact Hausdorff space. A continuous function $f : X \rightarrow \mathbb{C}$ *vanishes at infinity* if, for each $\epsilon > 0$, there is a compact set $C_\epsilon \subseteq X$ such that $X \setminus C_\epsilon \subseteq \{x \in X : |f(x)| < \epsilon\}$. Let $C_0(X)$ denote the Banach space of all functions on X which vanish at infinity, equipped with the norm $\|f\|_\infty := \sup_{x \in X} |f(x)|$. The space $C_c(X)$ of continuous functions with compact support is norm dense in $C_0(X)$ [3, III.1, Exercise 13]; when X is compact, $C_0(X) = C_c(X) = C(X)$, where $C(X)$ denotes the space of continuous complex-valued functions on X . The Riesz Representation Theorem [3, C.18] states that $C_0(X)^*$, the dual space of $C_0(X)$, is isometrically isomorphic to $M(X)$, the space of finite regular complex Borel measures on X (equipped with the norm $\|\mu\| := |\mu(X)|$); under this duality, corresponding to $\mu \in M(X)$ is the functional $\hat{\mu}$ on $C_0(X)$ defined by $\hat{\mu}(f) := \int f d\mu$.

We now focus on the case where μ is a positive Borel measure on \mathbb{C}^d with convergent moments up to (at least) order n , i.e., $\mathbb{C}_{n,d}[z, \bar{z}] \subseteq L^1(\mu)$. Let $K := \text{supp } \mu$; without loss of generality, in the sequel we normalize μ so that $\mu(K) = 1$. Since the monomials $\bar{z}^i z^j$ ($|i| + |j| \leq n$) are absolutely

integrable, we may consider the corresponding moments of μ ,

$$\gamma_{ij} := \int_K \bar{z}^i z^j d\mu(z), \quad |i| + |j| \leq n.$$

Let $\rho_0(z) := \|z\|^n$ (where, as usual, $\|z\| := (|z_1|^2 + \dots + |z_d|^2)^{1/2}$). Let $\Gamma := \int_K \rho_0(z) d\mu(z)$. If n is even, say $n = 2m$, then $\|z\|^n = (\bar{z}_1 z_1 + \dots + \bar{z}_d z_d)^m \in \mathbb{C}_{n,d}[z, \bar{z}]$, so $\Gamma < +\infty$. For the case when n is odd, to see that $\Gamma < +\infty$ note that by the equivalence of all norms on \mathbb{C}^d , there exists a constant $M > 0$ such that for every $z \in \mathbb{C}^d$, $\|z\| \leq M(|z_1| + \dots + |z_d|)$. Then $\int_K \|z\|^n d\mu(z) \leq M^n \int_K (|z_1| + \dots + |z_d|)^n d\mu(z)$, and the latter integral is convergent, since for every multi-index i with $|i| = n$, $z^i (= z_1^{i_1} \cdot \dots \cdot z_d^{i_d})$ is absolutely integrable.

Let $\gamma \equiv \gamma^{(n)}[\mu] := \{\gamma_{i,j}\}_{0 \leq |i|+|j| \leq n}$, and set $V(\mu; n) := \{\nu \in M(K) : \nu \geq 0, \gamma_{ij} = \int_K \bar{z}^i z^j d\nu(z), |i| + |j| \leq n-1, \int_K \|z\|^n d\nu(z) \leq \Gamma\}$. Observe that $V(\mu; n)$ is convex, and is nonempty since $\mu \in V(\mu; n)$. For $\nu \in V(\mu; n)$, $\|\nu\| = \nu(K) = \gamma_{00} = 1$, so $V(\mu; n)$ embeds as a subset of $B_1(C_0(K)^*)$, the closed unit ball of $C_0(K)^*$; recall that $B_1(C_0(K)^*)$ is weak-* compact and metrizable [3, Theorems V.3.1 and V.5.1].

PROPOSITION 3.2. *$V(\mu; n)$ is weak-* closed in $B_1(C_0(K)^*)$.*

To prove Proposition 3.2 we rely on the following technical result of Stochel.

PROPOSITION 3.3. *([22, Proposition 1]) Let F be a nonempty closed subset of \mathbb{C}^d and let ρ be a non-negative continuous function on F . Assume that $\{\mu_w\}_{w \in \Omega}$ is a net of finite positive Borel measures on F and μ is a finite positive Borel measure on F such that*

$$(i) \lim_{w \in \Omega} \int_F f d\mu_w = \int_F f d\mu \quad (f \in C_c(F)), \text{ and}$$

$$(ii) \sup_{w \in \Omega} \int_F \rho d\mu_w < +\infty.$$

Then $\int_F \rho d\mu \leq \sup_{w \in \Omega} \int_F \rho d\mu_w$, and $\int_F f \rho d\mu = \lim_{w \in \Omega} \int f \rho d\mu_w$ ($f \in C_0(F)$). Moreover, if the set $\{z \in F : \rho(z) \leq r\}$ is compact for some $r > 0$, then $\int_F f d\mu = \lim_{w \in \Omega} \int_F f d\mu_w$ for every $f : F \rightarrow \mathbb{C}$ such that $\frac{f}{1+\rho} \in C_0(F)$.

(The estimate on $\int_F \rho d\mu$ is not part of the statement of [22, Proposition 1], but is established in the proof.)

Proof. of Proposition 3.2. Since $B_1(C_0(K)^*)$ is weak-* metrizable, to establish that $V(\mu; n)$ is weak-* closed, it suffices to show that it is closed

under limits of sequences. Let $\{\nu_k\}_{k=1}^\infty \subseteq V(\mu; n)$ and suppose $\{\nu_k\}$ is weak-* convergent to $\Lambda \in B_1(C_0(K)^*)$, i.e., $\Lambda(f) = \lim_{k \rightarrow \infty} \int f d\nu_k$ ($f \in C_0(K)$). Clearly, $\Lambda \geq 0$, since each $\nu_k \geq 0$. Thus, by the Riesz Representation Theorem, there exists a finite, positive, regular Borel measure ν , $\text{supp } \nu \subseteq K$, such that $\Lambda = \hat{\nu}$.

We claim that $\nu \in V(\mu; n)$. Let $\rho = \rho_0$, i.e., $\rho(z) := \|z\|^n$ ($z \in K$). Since $\nu_k \in V(\mu; n)$, then $\int_K \rho(z) d\nu_k(z) \leq \Gamma$ ($\equiv \int_K \rho(z) d\mu$), whence Proposition 3.3 implies $\int_K \rho(z) d\nu \leq \Gamma$. To complete the proof, we will show that for $|i| + |j| \leq n - 1$, $f_{ij}(z, \bar{z}) := \bar{z}^i z^j$ satisfies

$$\int_K f_{ij} d\nu(z) = \gamma_{ij}. \quad (3.1)$$

For each k , $\int_K f_{ij} d\nu_k(z) = \gamma_{ij}$; Proposition 3.3 implies that to establish (3.1) it suffices to verify that $\frac{f_{ij}}{1+\rho} \in C_0(K)$ ($|i| + |j| \leq n - 1$). Let $L > 1$ and suppose $\|z\|^2 > L^2$, i.e., z is in the complement of the compact set $\{z \in \mathbb{C}^d : \|z\| \leq L\}$. Choose $i(z)$, $1 \leq i(z) \leq d$, such that $|z_{i(z)}| \geq |z_i|$, $1 \leq i \leq d$ ($i(z)$ depends on z). Then

$$|z_{i(z)}|^2 \geq \frac{\|z\|^2}{d} > \frac{L^2}{d},$$

so

$$\frac{1}{|z_{i(z)}|} < \frac{\sqrt{d}}{L}.$$

Now

$$\begin{aligned} \frac{|f_{ij}|}{1+\rho} &= \frac{|z_1|^{i_1+j_1} \dots |z_d|^{i_d+j_d}}{1 + \left(|z_1|^2 + \dots + |z_d|^2\right)^{n/2}} \leq \frac{|z_{i(z)}|^{|i|+|j|}}{|z_{i(z)}|^n} \\ &= \frac{1}{|z_{i(z)}|^{n-|i|-|j|}} < \frac{d^n}{L} \longrightarrow 0 \quad (L \longrightarrow +\infty). \end{aligned}$$

Thus $\frac{f_{ij}}{1+\rho} \in C_0(K)$ and the proof is complete. ■

We next require a variant of Proposition 2.2.

LEMMA 3.4. *If ν is an extreme point of $V(\mu; n)$, then ν is finitely atomic, with $\text{card supp } \nu \leq 1 + \dim \mathbb{C}_{n-1,d}[z, \bar{z}](\nu)$.*

Proof. Since ν is a representing measure for $\{\gamma_{ij}\}_{|i|+|j| \leq n-1}$, and $\int \|z\|^n d\nu < +\infty$, we may consider the subspace of $L^1(\nu)$ defined by

$\mathcal{M} := \{[p] + \alpha[\rho_0] : p \in \mathbb{C}_{n-1,d}[z, \bar{z}], \alpha \in \mathbb{C}\}$. We claim that \mathcal{M} is dense in $L^1(\nu)$. Since $\rho_0 \geq 0$, \mathcal{M} is self-adjoint; thus, if \mathcal{M} is not dense, it follows as in the proof of Proposition 2.2 that there exists $f : \text{supp } \nu \rightarrow \mathbb{R}$, $\|f\|_\infty \leq \frac{1}{2}$, such that $\int pf \, d\nu = 0$ for every $p \in \mathcal{M}$. Let $\nu_1 := (1+f)\nu$ and $\nu_2 := (1-f)\nu$; for $i = 1, 2, \nu_i \geq 0$, ν_i is a representing measure for $\{\gamma_{ij}\}_{|i|+|j| \leq n-1}$, and

$$\int \|z\|^n \, d\nu_i = \int \|z\|^n \, d\nu \pm \int \|z\|^n f \, d\nu = \int \|z\|^n \, d\nu \leq \Gamma.$$

Thus $\nu_i \in V(\mu; n)$ ($i = 1, 2$), and since $\nu = \frac{1}{2}\nu_1 + \frac{1}{2}\nu_2$, we have a contradiction to the hypothesis that ν is an extreme point. The rest of the proof is identical to that of Proposition 2.2; in particular, since \mathcal{M} is finite dimensional,

$$\text{card supp } \nu \leq \dim L^\infty(\nu) = \dim L^1(\nu) = \dim \mathcal{M} \leq 1 + \dim \mathbb{C}_{n-1,d}[z, \bar{z}](\nu).$$

■

Proof. of Theorem 3.1. Since $\mu \in V(\mu; n)$, Proposition 3.2 implies that $V(\mu; n)$ is a nonempty weak-* compact convex subset of $B_1(C_0(K)^*)$ (where $K := \text{supp } \mu$). It follows from the Krein-Millman Theorem that $V(\mu; n)$ has an extreme point, and Lemma 3.4 implies that any such extreme point ν corresponds to a quadrature rule for μ of degree $n-1$ with size at most $1 + N_{n-1,d}(\nu) \leq 1 + N_{n-1,d;\nu} \leq 1 + N_{n-1,d;\mu}$. ■

We next turn to the complex version of Theorem 1.1.

THEOREM 3.5. *Suppose μ is a positive finite Borel measure on \mathbb{C}^d with compact support K . Given $m > 0$, there exist $N \leq \dim(\mathbb{C}_{m,d}[z, \bar{z}]|_K)$, nodes z_1, \dots, z_N in K , and positive weights ρ_1, \dots, ρ_N such that*

$$\int_K p(z, \bar{z}) \, d\mu(z) = \sum_{i=1}^N \rho_i p(z_i, \bar{z}_i) \quad (p \in \mathbb{C}_{m,d}[z, \bar{z}]).$$

Proof. Since $C_0(K) = C(K)$, it is straightforward to modify the proof of Proposition 3.2 to conclude that $M_K(\gamma^{(m)}[\mu])$ is weak-* closed in $B_1(C_0(K)^*)$. Thus $M_K(\gamma^{(m)}[\mu])$ has an extreme point ν , and Proposition 2.2 implies $\text{card supp } \nu \leq \dim \mathbb{C}_{m,d}[z, \bar{z}]|_{\text{supp } \nu} \leq \dim \mathbb{C}_{m,d}[z, \bar{z}]|_K$. ■

Proof. of Theorems 1.1, 1.3 and 1.4. The proofs of Theorems 3.1 and 3.5 depend on duality results for $L^1(\mu)^*$ and $C_0(K)^*$, where μ is a positive Borel measure on \mathbb{C}^d and K is a closed subset of \mathbb{C}^d . These duality results,

including those of Stochel (e.g., Proposition 3.3), admit exact analogues for the real case, where \mathbb{C}^d is replaced by \mathbb{R}^d . Thus we may formally repeat the proof of Theorem 3.1 to obtain Theorem 1.4 (and Theorem 1.3), and similarly for Theorem 3.5 and Theorem 1.1. (For the case when $d = 2p$, alternate proofs can be based on the equivalence of moment problems on \mathbb{R}^{2p} with those on \mathbb{C}^p .) ■

Implicit in [25, pp. 127-129] is a representation theorem for any positive linear functional on $\mathbb{R}_m[t] \upharpoonright_K$ ($K \subseteq \mathbb{R}^d$ compact). We next formulate this result for the complex case and give a new proof, based not on convexity but on C^* -algebra ideas.

PROPOSITION 3.6. *(cf. [25, Théorème II]) Let $m > 0$ and let K be a compact subset of \mathbb{C}^d . If $\Phi : \mathbb{C}_{m,d}[z, \bar{z}] \upharpoonright_K \rightarrow \mathbb{C}$ is a positive linear functional, then there exist $N \leq \dim \mathbb{C}_{m,d}[z, \bar{z}] \upharpoonright_K$, nodes z_1, \dots, z_N in K , and positive weights ρ_1, \dots, ρ_N , such that $\Phi(p) = \sum_{i=1}^N \rho_i p(z_i, \bar{z}_i)$ ($p \in \mathbb{C}_{m,d}[z, \bar{z}] \upharpoonright_K$).*

Proof. $\mathbb{C}_{m,d}[z, \bar{z}]$ is an operator system in $C(K)$ (cf. [16, Chapter 2] and [4, Chapter 5, Definition 33.1]), so Φ can be extended to a positive linear functional $\tilde{\Phi}$ on $C(K)$ ([16, Exercice 2.10] or [4, Chapter 5, Proposition 33.2(c)]). By the Riesz Representation Theorem, there is a positive Borel measure μ , $\text{supp } \mu \subseteq K$, such that $\tilde{\Phi}(f) = \int_K f d\mu$ ($f \in C(K)$). The result now follows by applying Theorem 3.5 to μ . ■

4. UNIQUE REPRESENTING MEASURES IN TCMP

Concerning the analogue of Question 1.6 for the truncated complex moment problem, consider the following possible properties of $\gamma \equiv \gamma^{(2n)}$:

- (P1) γ has a unique representing measure;
- (P2) $M(\gamma) := \{\nu : \nu \text{ is a representing measure for } \gamma\}$ is weak-* closed in $B_1(C_0(\mathbb{C})^*)$;
- (P3) γ admits a finitely atomic representing measure.

Results of the preceding sections show that (P1) \Rightarrow (P2) \Rightarrow (P3). In the sequel we establish (P1) in two basic cases of TCMP.

Recall that γ is flat if $M(n) \equiv M(n)(\gamma) \geq 0$ and $\text{rank } M(n) = \text{rank } M(n-1)$. In [5, Corollary 5.14] we proved that if γ is flat, then γ admits a unique representing measure having moments of all orders (and this measure is rank $M(n)$ -atomic).

PROPOSITION 4.1. *If $\gamma^{(2n)}$ is flat, then there exists a unique representing measure, which is rank $M(n)$ -atomic.*

Proof. Let μ be a representing measure for $\gamma^{(2n)}$. Since [5, Corollary 5.14] implies that γ has a unique representing measure having moments of all orders, it suffices to establish that μ has moments of all orders. We first consider moments of order $2n + 1$. Suppose $i, j \geq 0$ and $i + j = n$; since $M(n) \geq 0$ and $\text{rank } M(n) = \text{rank } M(n - 1)$, there exists $p_{ij} \in \mathbb{C}_{n-1}[z, \bar{z}]$ such that $\bar{Z}^i Z^j = p_{ij}(Z, \bar{Z}) \in \mathcal{C}_{M(n)}$ (the column space of $M(n)$). Since μ is a representing measure, $\bar{z}^i z^j = p_{ij}(z, \bar{z})$ on $\text{supp } \mu$ [5, Proposition 3.1]. Thus, for $k, l \geq 0$, $k + l = n + 1$, $\bar{z}^{i+k} z^{j+l} = (\bar{z}^k z^l p_{ij})(z, \bar{z})$ on $\text{supp } \mu$, and since $\deg \bar{z}^k z^l p_{ij} \leq 2n$, then $\int \bar{z}^{i+k} z^{j+l} d\mu$ is convergent. By considering all indices i, j, k, l with $i + j = n$ and $k + l = n + 1$, it follows that μ has convergent moments up to degree $2n + 1$.

We next consider degree $2n + 2$. There exists $p_{0,n} \in \mathbb{C}_{n-1}[z, \bar{z}]$ such that $Z^n = p_{0,n}(Z, \bar{Z})$, whence $z^n = p_{0,n}(z, \bar{z})$ on $\text{supp } \mu$. Thus $|z|^{2n+2} = (|z|^2 |p_{0,n}|^2)(z, \bar{z})$ on $\text{supp } \mu$; now $\deg |z p_{0,n}|^2 \leq 2n$, so $\int |z|^{2n+2} d\mu < +\infty$. Thus, for all (i, j) such that $i + j = 2n + 2$, $\int \bar{z}^i z^j d\mu$ is absolutely convergent, hence convergent.

Since μ has convergent moments up to degree $2n + 2$, we may consider $M(n + 1)[\mu]$. Since μ is a representing measure for $M(n + 1)[\mu]$, it follows that $M(n + 1)[\mu]$ is positive and recursively generated. For $i + j = n$, we have $\bar{Z}^i Z^j = p_{ij}(Z, \bar{Z})$ in $\mathcal{C}_{M(n)}$, and since $M(n + 1)[\mu] \geq 0$, it follows that $\bar{Z}^i Z^j = p_{ij}(Z, \bar{Z})$ in $\mathcal{C}_{M(n+1)[\mu]}$ [10, Proposition 2.4 (Extension Principle)]. By recursiveness, we have $\bar{Z}^i Z^{j+1} = (z p_{ij})(Z, \bar{Z})$ and $\bar{Z}^{i+1} Z^j = (\bar{z} p_{ij})(Z, \bar{Z})$ in $\mathcal{C}_{M(n+1)[\mu]}$. Since $\deg z p_{ij} \leq n$ and $\deg \bar{z} p_{ij} \leq n$, it follows that for $k + l = n + 1$, $\bar{Z}^k Z^l \in \langle \bar{Z}^i Z^j \rangle_{0 \leq i+j \leq n}$ in $\mathcal{C}_{M(n+1)[\mu]}$, whence $\text{rank } M(n + 1)[\mu] = \text{rank } M(n)[\mu]$.

Thus, $\gamma^{2(n+1)}[\mu]$ is flat. The preceding argument may be repeated to produce successive flat extensions $M(n + 1)[\mu], M(n + 2)[\mu], \dots$; thus μ has convergent moments of all orders. ■

PROPOSITION 4.2. *If $\gamma^{(2n)}$ has a representing measure and if $M(n)(\gamma)$ admits an analytic relation as in (1.3) for some $k \leq n$, then γ has a unique representing measure, which is finitely atomic with at most k^2 atoms.*

REMARK 4.3. [12, Theorem 3.1] illustrates the case of $\gamma \equiv \gamma^{(6)}$ such that $M(3)(\gamma)$ has an analytic relation with $k = n = 3$, and such that the unique representing measure is 9-atomic.

We require the following preliminary result, which is of independent interest.

PROPOSITION 4.4. *A polynomial of the form $p(z, \bar{z}) \equiv z^k - q(z, \bar{z})$, where $q \in \mathbb{C}_{k-1}[z, \bar{z}]$, has at most k^2 roots.*

Proof. The result is obvious for $k = 1$, so we assume $k \geq 2$. Let $\mathcal{Z}(p)$ denote the zero set of p and assume that $\mathcal{Z}(p) \supseteq \Lambda \equiv \{z_1, \dots, z_m\}$, where $m := k^2 + 1$ and the z_i s are distinct. Let μ be a positive measure with $\text{supp } \mu = \Lambda$, let $n := 2k - 2$, and form $M(n)[\mu]$. Clearly μ is a representing measure for $M(n)[\mu]$, and since $\deg p \leq n$ and $\text{supp } \mu \subseteq \mathcal{Z}(p)$, we must have $p(Z, \bar{Z}) = 0$ in $\mathcal{C}_{M(n)[\mu]}$ ([5, Proposition 3.1]). It follows that $Z^k = q(Z, \bar{Z})$, with $k = \frac{n}{2} + 1 = \lceil \frac{n}{2} \rceil + 1$ and $\deg q < k$. By [6, Theorem 3.1], $M(n)[\mu]$ admits a unique positive, recursively generated extension $M \equiv M(\infty)$, and M is a flat extension. Since $M(\infty)[\mu]$ is positive and recursively generated, we have $M(\infty)[\mu] = M$, and [5, Proposition 4.6] implies that

$$m = \text{card supp } \mu = \text{rank } M(\infty)[\mu] = \text{rank } M = \text{rank } M(n)[\mu].$$

Observe now that the size of $M(n)[\mu]$ is $\frac{(n+1)(n+2)}{2} = k(2k-1)$. The column relation $Z^k = q(Z, \bar{Z})$ (and the conjugate relation $\bar{Z}^k = \bar{q}(Z, \bar{Z})$) propagate recursively (and disjointly) in $\mathcal{C}_{M(n)[\mu]}$ to generate $2(1 + 2 + \dots + k - 1) = k(k-1)$ columns each of which is a linear combination of columns of strictly lower degree. (These $k(k-1)$ columns are $Z^k, \bar{Z}^k, Z^{k+1}, \bar{Z}Z^k, \dots, \bar{Z}^k Z, \bar{Z}^{k+1}, \dots, Z^n, \dots, \bar{Z}^{n-k} Z^k, \bar{Z}^k Z^{n-k}, \dots, \bar{Z}^n$.) Thus,

$$m = \text{rank } M(n)[\mu] \leq k(2k-1) - k(k-1) = k^2 < m,$$

a contradiction. We conclude that $\text{card } \mathcal{Z}(p) \leq k^2$. ■

REMARK 4.5. *The following examples show that the estimate in Proposition 4.4 is sharp. Let $p(z, \bar{z}) := z^2 - \bar{z}$. Then $\mathcal{Z}(p) = \{0\} \cup \{\omega^j : 0 \leq j \leq 2\}$, where ω is a primitive cubic root of unity. Thus, $\text{card } \mathcal{Z}(p) = 4 = (\deg p)^2$. For $k = 3, 4, 5$, A. Wilmschurst [26, Chapter 3, Section 5] has constructed polynomials $q_3(z, \bar{z}) := z^3 + z^2 + z - 2\bar{z}^2 - 2\bar{z}$, $q_4(z, \bar{z}) := z^4 + 3z^2 - z - 3\bar{z}^3 - 3\bar{z}^2$ and $q_5(z, \bar{z}) := z^5 + 5z^3 - 10z^2 + 5z + 5\bar{z}^4 - 5\bar{z}^3$, with $\text{card } \mathcal{Z}(q_k) = (\deg q_k)^2$ ($k = 3, 4, 5$).*

Proof of Proposition 4.2. We have an analytic relation of the form $Z^k = q(Z, \bar{Z})$ for some $k \leq n$, where $q \in \mathbb{C}_{k-1}[z, \bar{z}]$. Let $p(z, \bar{z}) := z^k - q(z, \bar{z})$; Proposition 4.4 implies that $\text{card } \mathcal{Z}(p) \leq k^2$. Let μ be a representing measure for γ . Since $\text{supp } \mu \subseteq \mathcal{Z}(p)$ [5, Proposition 3.1], $\text{card supp } \mu \leq k^2$. Thus μ has moments of all orders, and recursiveness implies that $M \equiv M(\infty)[\mu]$ is completely determined by the analytic column relations

$$Z^{k+j} = (z^j q)(Z, \bar{Z}) \quad (j \geq 0). \quad (4.4.1)$$

(Note that for each p , $M(p+1)[\mu]$ is completely determined by $M(p)[\mu]$ and the moments in Z^{p+1} .) Further, since μ is a representing measure for M , $\text{rank } M \leq \text{card supp } \mu \leq k^2$ [5, Proposition 4.6]. Now, if ν is a representing measure for γ , then it follows as above that $M(\infty)[\nu]$ is also determined by (4.4.1), whence $M(\infty)[\nu] = M$. Since M is positive and has finite rank, [5, Theorem 4.7] implies that it has a unique representing measure, whence $\mu =$

ν . ■

Added in Proof. We have recently learned from Professor Christian Berg that the weak-* convergence techniques used in the proof of Theorem 1.4 have antecedents in the literature; a reference for these techniques is [2, Chapter 2].

REFERENCES

1. N.I. Akhiezer, "The Classical Moment Problem," Hafner Publ. Co., New York, 1965.
2. C. Berg, J.P.R. Christensen and P. Ressel, "Harmonic Analysis on Semigroups," Springer-Verlag, New York, 1984.
3. J. Conway, "A Course in Functional Analysis," Graduate Texts in Mathematics, Springer-Verlag, New York, 1990.
4. J. Conway, "A Course in Operator Theory," Graduate Studies in Mathematics, Vol. 21, American Mathematical Society, Providence, 2000.
5. R. Curto and L. Fialkow, Solution of the truncated complex moment problem for flat data, *Mem. Amer. Math. Soc.* 119 (1996), no. 568.
6. R. Curto and L. Fialkow, Flat extensions of positive moment matrices: Relations in analytic or conjugate terms, *Oper. Theory Adv. Appl.* 104 (1998), 59–82.
7. R. Curto and L. Fialkow, Flat extensions of positive moment matrices: Recursively generated relations, *Mem. Amer. Math. Soc.* 136 (1998), no. 648.
8. R. Curto and L. Fialkow, The truncated complex K -moment problem, *Trans. Amer. Math. Soc.* 352 (2000), 2825–2855.
9. R. Curto and L. Fialkow, Solution of the singular quartic moment problem, *J. Operator Theory*, to appear.
10. L. Fialkow, Positivity, extensions, and the truncated complex moment problem, in "Multivariable operator theory (Seattle, WA, 1993)," *Contemp. Math.*, vol. 185, American Mathematical Society, Providence, 1995, pp. 133–150.
11. L. Fialkow, Multivariable quadrature and extensions of moment matrices, preprint 1996.
12. L. Fialkow, Minimal representing measures arising from rank-increasing moment matrix extensions, *J. Operator Theory* 42 (1999), 425–436.
13. J.P. Gabardo, Truncated trigonometric moment problems and determinate measures, preprint 1999.
14. H.M. Möller, Kubaturformeln mit minimaler Knotenzahl, *Numer. Math.* 25 (1976), 185–200.
15. I.P. Mysovskikh, On Chakalov's theorem, *U.S.S.R. Comput. Math. and Math. Phys.* 15 (1975), no. 6, 221–227; (translation in *Zh. Vychisl. Mat. i Mat. Fiz.* 15 (1975), no. 6, 1589–1593, 1627).
16. V. Paulsen, "Completely bounded maps and dilations," *Pitman Res. Notes Math. Ser.*, vol. 146, Longman Sci. Tech., London, 1986.
17. M. Putinar, On Tchakaloff's Theorem, *Proc. Amer. Math. Soc.* 125 (1997), 2409–2414.
18. A. Ralston, "A First Course in Numerical Analysis," McGraw-Hill, New York, 1965.
19. B. Reznick, Sums of even powers of real linear forms, *Mem. Amer. Math. Soc.* 96 (1992), no. 463.
20. H. Schmid, Two-dimensional minimal cubature formulas and matrix equations, *SIAM J. Matrix Anal. Appl.* 16(1995), 898–921.
21. H. Schmid and Y. Xu, On bivariate Gaussian cubature formulae, *Proc. Amer. Math. Soc.* 122(1994), 833–841.
22. J. Stochel, Solving the truncated moment problem solves the full moment problem, *Glasgow Math. J.*, to appear.

23. J. Stochel and F. Szafraniec, The complex moment problem and subnormality: A polar decomposition approach, *J. Funct. Anal.* 159(1998), 432-491.
24. A.H. Stroud, "Approximate Calculation of Multiple Integrals," Prentice-Hall, Englewood Cliffs, N.J., 1971.
25. V. Tchakaloff, Formules de cubatures mécaniques à coefficients non négatifs, *Bull. Sci. Math. (2)* 81 (1957), 123-134.
26. A. Wilmschurst, Complex Harmonic Mappings and the Valence of Harmonic Polynomials, Ph.D. Dissertation, University of York, 1994.
27. Y. Xu, "Common zeros of polynomials in several variables and higher-dimensional quadrature," Pitman Res. Notes Math. Ser., vol. 312, Longman Sci. Tech., Harlow, 1994.