

Haagerup-Schultz projections, upper triangular forms and spectrality of Dixmier traces

Ken Dykema

Department of Mathematics
Texas A&M University
College Station, TX, USA.

University of Iowa
Workshop on Noncommutative Analysis
June 4, 2016

(this talk was given on the whiteboard)

Selected References (chronological)

- [1] Brown, *Lidskii's theorem in the type II case*, Geometric methods in operator algebras (Kyoto, 1983), Pitman Res. Notes Math. Ser., vol. 123, Longman Sci. Tech., Harlow, 1986, pp. 1–35.
- [2] Haagerup and Schultz, *Brown measures of unbounded operators affiliated with a finite von Neumann algebra*, Math. Scand. **100** (2007), 209–263.
- [3] ———, *Invariant subspaces for operators in a general II_1 -factor*, Publ. Math. Inst. Hautes Études Sci. **109** (2009), 19–111.
- [4] Dykema, Sukochev, and Zanin, *A decomposition theorem in II_1 -factors*, J. reine angew. Math. **708** (2015), 97–114.
- [5] ———, *Holomorphic functional calculus on upper triangular forms in finite von Neumann algebras*, Illinois J. Math., to appear, available at [arXiv:1310.2524](https://arxiv.org/abs/1310.2524).
- [6] Noles, *Upper triangular forms and spectral orderings in a II_1 -factor*, available at [arXiv:1406.2774](https://arxiv.org/abs/1406.2774).
- [7] Dykema, Sukochev, and Zanin, *An upper triangular decomposition theorem for some unbounded operators affiliated to II_1 -factors*, available at [arXiv:1509.03358](https://arxiv.org/abs/1509.03358).
- [8] ———, *Algebras of log-integrable functions and operators*, Complex Anal. Oper. Theory, to appear, available at [arXiv:1509.03360](https://arxiv.org/abs/1509.03360).
- [9] Dykema, Noles, Sukochev, and Zanin, *On reduction theory and Brown measure for closed unbounded operators*, available at [arXiv:1509.03362](https://arxiv.org/abs/1509.03362).